Characterizing Air Temperature Changes in the Tarim Basin over 1960–2012



Dongmei Peng^{1,2,6}, Xiujun Wang^{1,3}*, Chenyi Zhao¹, Xingren Wu⁴, Fengqing Jiang¹, Pengxiang Chen⁵

1 State Key Laboratory of Desert and Oasis Ecology, Xinjiang Institute of Ecology and Geography, Chinese Academy of Sciences, Urumqi, Xinjiang, China, 2 Graduate University of Chinese Academy of Sciences, Beijing, China, 3 Earth System Science Interdisciplinary Center, University of Maryland, College Park, Maryland, United States of America, 4 IM System Group and Environmental Modeling Center, National Centers for Environmental Prediction, National Oceanic and Atmospheric Administration, College Park, Maryland, United States of America, 5 Xinjiang Meteorological Observatory, Urumqi, Xinjiang, China, 6 Information Centre of Xinjiang Xingnong-Net, Urumqi, China

Abstract

There has been evidence of warming rate varying largely over space and between seasons. However, little has been done to evaluate the spatial and temporal variability of air temperature in the Tarim Basin, northwest China. In this study, we collected daily air temperature from 19 meteorological stations for the period of 1960-2012, and analyzed annual mean temperature (AMT), the annual minimum (T_{min}) and maximum temperature (T_{max}), and mean temperatures of all twelve months and four seasons and their anomalies. Trend analyses, standard deviation of the detrended anomaly (SDDA) and correlations were carried out to characterize the spatial and temporal variability of various mean air temperatures. Our data showed that increasing trend was much greater in the T_{min} (0.55°C/10a) than in the AMT (0.25°C/10a) and T_{max} (0.12°C/10a), and the fluctuation followed the same order. There were large spatial variations in the increasing trends of both AMT (from -0.09 to 0.43 °C/10a) and T_{min} (from 0.15 to 1.12°C/10a). Correlation analyses indicated that AMT had a significantly linear relationship with T_{min} and the mean temperatures of four seasons. There were also pronounced changes in the monthly air temperature from November to March at decadal time scale. The seasonality (i.e., summer and winter difference) of air temperature was stronger during the period of 1960–1979 than over the recent three decades. Our preliminary analyses indicated that local environmental conditions (such as elevation) might be partly responsible for the spatial variability, and large scale climate phenomena might have influences on the temporal variability of air temperature in the Tarim Basin. In particular, there was a significant correlation between index of El Niño-Southern Oscillation (ENSO) and air temperature of May (P = 0.004), and between the index of Pacific Decadal Oscillation (PDO) and air temperature of July (P = 0.026) over the interannual to decadal time scales.

Citation: Peng D, Wang X, Zhao C, Wu X, Jiang F, et al. (2014) Characterizing Air Temperature Changes in the Tarim Basin over 1960–2012. PLoS ONE 9(11): e112231. doi:10.1371/journal.pone.0112231

Editor: Inés Álvarez, University of Vigo, Spain

Received June 7, 2014; Accepted October 2, 2014; Published November 6, 2014

Copyright: © 2014 Peng et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability: The authors confirm that all data underlying the findings are fully available without restriction. Data are available from the National Climatic Center, China Meteorological Administration Institutional Data Access/Ethics Committee. Requests to access data can be sent to: Mr. Pengxiang Chen Xinjiang Meteorological Observatory Urumqi 830002, Xinjiang, China Email: cpx1860@163.com

Funding: This research was supported by the Chinese National Basic Research Key Project (No. 2010cb951001), National Natural Science Foundation (No. 41171095), and National Science & Technology Pillar Program (No. 2013BAC10B01). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* Email: wwang@essic.umd.edu

Introduction

There have been numerous studies of temporal variations of air temperature at various spatial scales [1–5], which all have pointed to the fact that there has been a general warming trend in the global mean air temperature. Earlier studies indicated that the magnitude of warming in the Northern Hemisphere (0.30° C/10a) was more than double of the one (0.13° C/10a) in the Southern Hemisphere during 1977–2001 [6,7].

There has been evidence of difference in warming trend over space and time in China. For example, the warming rate of air temperature was 0.25° C/10a for the period of 1951–2004 in China [8], but 0.35° C/10a in northwest China during the period of 1961–2006 [9], which were much higher than the global average. Over the past 50 years, an increase of air temperature with a linear tendency of 0.28° C/10a was observed in Xinjiang, which was lower than that for the northwest China [10]. A couple

of studies showed that the warming in the Tarim Basin was only from 0.19 to 0.22° C/10a [11,12]. These findings indicated that climate change in Xinjiang might have its own spatial and temporal characteristics due to its large extent and complex terrain.

Apart from the increasing trend in annual mean temperature (AMT), there has been evidence of differences in warming rate between seasons at various regional scales. An earlier study based on analyses of 726 stations' data (1951–2004) in China showed an increasing trend of 0.39° C/10a in winter, 0.28° C/10a in spring, 0.20° C/10a in autumn and only 0.15° C/10a in summer [8]. Some studies indicated that the increasing rate of individual season's temperature was even greater in northwest China, in particular in winter, 0.35° C/10a in autumn, 0.26° C/10a in symple, Wang et al. [13] reported a rate of 0.56° C/10a in winter, 0.35° C/10a in autumn, 0.26° C/10a in spring, and 0.22° C/10a in summer for the period of 1960–2005. A



Figure 1. Topographic map with the locations of the meteorological stations in Tarim Basin and climatology AMT for the period of 1960–2012. doi:10.1371/journal.pone.0112231.g001

recent study by Li et al. [14] showed an increasing trend of 0.49, 0.36, 0.27 and 0.23°C/10a for winter, autumn, spring and summer, respectively, for the period of 1960–2010. These results imply that there are significant changes in the seasonality of air temperature in the vast arid/semi-arid regions. However, there were limited analyses for the Tarim Basin, which were based on data of 1958–2004 from six meteorological stations in the western section of the basin [15], showing an increasing trend of 0.4 and 0.3° C/10a in winter and autumn, respectively.

Large scale climate phenomena often have various impacts on local climate. For example, El Niño-Southern Oscillation (ENSO) has widespread influence on the climate of north America and east Asia. An early study indicated that ENSO had significant effects on precipitation and temperature in the basins in Xinjiang [16]. However, other studies suggested that ENSO had no significant effect on the annual temperature in the Tarim basin [15,17]. On the other hand, there was evidence of interannual to decadal variations in air temperature in the Tarim Basin [17–19]. However, little is done to evaluate the temporal variability of air temperature change and the underlying mechanisms.

While there have been some studies of the climate changes in the Tarim Basin [12,15,20], there is a lack of consistence in findings, primarily due to the difference in the studying period (thus difference in dataset). In addition, detailed analyses of the spatial and temporal variations in air temperature have been lacking for the Tarim Basin, which hampers our understanding of the climate change at local to regional scales. To address this issue, we carry out a detailed study, using the latest dataset covering the period of 1958–2012. Our approaches include anomalies, trend analyses, and fluctuation and correlation analyses. The objective of this study is to investigate the spatial and temporal variability in air temperature at seasonal to decadal time scales, and to explore the possible mechanisms responsible for these changes.

Materials and Methods

Description of study area

The Tarim Basin is located in the southern part of the Xinjiang Autonomous Region, Northwestern China, which is surrounded by high mountains (i.e., Tianshan Mountain and Kunlun Mountain) (Figure 1). The main river system, the Tarim River, consists of the Yarkant River, Aksu River, Hotan River, and Kaidu River. Rivers are primarily fed by snow and glacier melting waters from the mountains. The Tarim Basin (with a total area of 1.02×10^6 km²) is divided into the mountains (47%, with elevation of 4000–6000 m), plains (22%, with elevation of 800–1400 m), and deserts (31%, with elevation of 1200–1500 m in the western and southern parts, and 800–1000 m in the eastern and northern parts). The climate is typical continental, with an average annual temperature of 10.6–11.5°C and precipitation of 17.4–42.0 mm [21].

Data sources and analyses

Data sources. We obtained daily mean air temperature data of 19 meteorological stations for the period of 1960–2012 from the National Climatic Center, China Meteorological Administration (http://cmdp.ncc.cma.gov.cn/cn/index.htm). These stations have the most completed dataset with acceptable quality control [22]. Figure 1 shows the spatial distribution of these stations and AMT over the period of 1960–2012. As shown in Figure 1, AMT varies from 6.57°C to 12.75°C, with higher values in the southwest stations than in the northeast stations. However, the lowest AMT was found at the stations with higher elevation in the northwest section.

We obtained three common indices of large-scale climate variability, i.e., the Southern Oscillation Index (SOI), the Pacific Decadal Oscillation (PDO), and the Indian Ocean Dipole Mode Index (DMI). The SOI data were obtained from the Climate



Figure 2. Time series and linear trends of AMT and temperature anomalies in the Tarim Basin. Area averaged (a) AMT, and temperature anomaly for (b) annual minimum (solid line), and maximum (dotted line), (c) spring, (d) summer, (e) autumn, and (f) winter. doi:10.1371/journal.pone.0112231.g002

Prediction Center (CPC) of the National Weather Service, U.S. (http://www.cpc.ncep.noaa.gov/), PDO index from N. Mantua (http://jisao.washington.edu/pdo/), and DMI from the web site (http://www.jamstec.go.jp/frcgc/research/d1/iod/).

Data analyses. We used daily mean temperature to derive the minimum (T_{min}) and maximum (T_{max}) temperature, and average temperatures for all twelve months, spring (March-May), summer (June-August), autumn (September-November) and winter (December-February) for each year. Our analyses included anomalies, trend analysis, fluctuation analysis, and correlation. All calculations were performed using the SPSS 19.

Temperature anomaly (TA) was determined by removing the mean value (\overline{T}) that was calculated for the period of 1960–2012:

$$TA(x) = T_x - \overline{T}_x \tag{1}$$

where x represented any temperature index (e.g., T_{min} and T_{max}).

Trend analysis was carried out by linear regression that has been a common method to determine the long-term changing trend of air temperature [23]:

$$T^r = T_0^r + S(t - 1960) \tag{2}$$

where T^r and T'_0 were the predicted temperature for year t and 1960, respectively. The slope S represented the trend of temperature increase, and the strength and significance of the linear increase were indicated by the regression coefficient (R) and P value. In general, when the P value was smaller than 0.05, the trend was significant.

Detrended fluctuation analysis (DFA) has proven useful in revealing the extent of long range correlations in time series [24,25]. We carried out a similar analysis but with a single time window to determine the extent of fluctuation in various temperature indices, which was the standard deviation of the detrended time series of anomaly (SDDA) and calculated as:

| | АМТ | T _{max} | T _{min} | T _{spr} | T _{sum} | T _{aut} | T _{win} |
|--|---|-------------------------------|------------------|------------------|------------------|------------------|------------------|
| Trend (°C/10a) | 0.25 | 0.12 | 0.55 | 0.23 | 0.17 | 0.27 | 0.36 |
| ж | 0.668** | 0.200 | 0.328* | 0.430** | 0.454** | 0.565** | 0.386** |
| SDDA (°C/10a) | 0.43 | 06.0 | 2.43 | 0.74 | 0.52 | 0.61 | 1.26 |
| R with AMT | | 0.259 | 0.533** | 0.564** | 0.616** | 0.701** | 0.335* |
| Significance of the regression/correlation wide to the second of the second signation wide to the second se | as marked with one (P $\scriptscriptstyle{	extsf{	imes}}$ | < 0.05) and two (P < 0.01) |) asterisks. | | | | |

$$SDDA = \sqrt{\frac{\sum_{i=1}^{n} (T_i - T_0^r - iS)^2}{n}}$$
 (3)

where T_i was the *i*th temperature, and *n* the number of data points. In general, the smaller the SDDA value, the smaller the fluctuation is.

Correlation analyses were carried out to examine the relationships between AMT and other temperatures and the potential impacts of climate phenomena over the period of 1961–2013. We used annual means of PDO and DMI indices and the mean value of SOI over winter. We evaluated the relationship between various temperature anomalies with ENSO, DMI and the PDO indices. We also conducted linear and nonlinear regression analyses to examine the relationship between the change rates of AMT or T_{min} and elevation, latitude and longitude. In addition, multiregressions were carried out to evaluate the influences of local factors and climate phenomena.

Correlation coefficient (R) was used to indicate the strength and direction of the relationship between the observation (x) and prediction (y), which was defined as the covariance of the variables divided by the product of their standard deviations:

$$R = \frac{n \sum xy - (\sum x)(\sum y)}{\sqrt{n(\sum x^2) - (\sum x)^2} \sqrt{n(\sum y^2) - (\sum y)^2}}$$
(4)

where *n* was the number of pairs of data. The value of R ranged from -1 to +1. The + and - signs indicated positive and negative linear correlations, respectively. The same as the regression analysis, the correlation coefficient *R* was assessed by the *P* value. When *P* was smaller than 0.05, the two variables were considered to have a significant correlation.

Results

Temperature change trend and fluctuation

We first evaluated the increasing trend of air temperature averaged over the entire basin. There was a significantly linear increase in the AMT (0.25°C/10a, P <0.001) and T_{min} (0.55°C/ 10a, P = 0.017) (Figure 2, Table 1). While the increasing trend of T_{max} (0.12°C/10a) was not significant, and weaker than that of $T_{\rm min},$ SDDA was smaller for $T_{\rm max}~(0.9)$ than for $T_{\rm min}~(2.43)$ (Table 1), indicating less fluctuation in T_{max} than in T_{min} relative to their change trends. Trend analyses also showed that there was a significantly (P < 0.05) linear increase in the mean temperatures of all seasons, with the greatest in winter $(0.36^{\circ}C/10a)$, followed by autumn ($0.27^{\circ}C/10a$), spring ($0.23^{\circ}C/10a$) and summer ($0.17^{\circ}C/$ 10a). However, comparisons of regressions indicated that these trends were not significantly different (P > 0.05, data not shown). SDDA was much smaller in summer than in other seasons, which indicated that temperature fluctuation relative to the trend was smaller in summer than in autumn, winter and spring in the Tarim Basin. Correlation analyses indicated that AMT had a significant relationship (P $<\!\!0.001\!)$ with $T_{\rm min}$ and all seasons' mean temperatures, but no significant relationship with T_{max} (R = 0.26, P = 0.061) (Table 1).

There was a considerably spatial variability in the increasing rate of AMT, ranging from -0.09° C/10a to 0.43° C/10a (Figure 3a), with only one station (i.e., N4) showing a decreasing



Figure 3. Spatial distributions of linear trend of AMT and its SDDA during 1960–2012 in the Tarim Basin. Spatial distributions of (a) linear trend of AMT (°C/10a) and (b) SDDA (°C/10a) of AMT. doi:10.1371/journal.pone.0112231.g003

trend (i.e., -0.09° C/10a). While there seemed no clear spatial pattern, the warming rate of AMT was similar (~0.26°C/10a) at all the stations located west of 79°E and east of the 86°E. The highest and lowest warming rates were found between 80°E and 86°E. Unlike the warming rate of AMT, the SDDA of AMT revealed a large spatial variation, with much higher values towards the west. In general, SDDA was larger at higher elevation stations than lower elevation stations, indicating that there would be less fluctuation in AMT in the regions with lower elevations.

Figure 4 shows the spatial pattern of the increasing rate of T_{min} , during 1960–2012. There was a big range in the increase of T_{min} , i.e., from 0.15 to 1.12°C/10a. In most parts of the Tarim Basin, T_{min} increasing rate reached 0.55°C/10a, which was almost twice as large as the rate of AMT. Clearly, T_{min} increasing rate was significantly higher in the west than in the east. In general, T_{min} revealed a greater increasing rate to the north than to the south, with very high rates (0.80°C/10a) in the most northern edge of the basin. Overall, the SDDA of T_{min} showed a similar spatial pattern to the increasing rate of T_{min} , i.e., higher values in the northern parts than in the southern parts, and to the west than to the east of the basin. The exception was at the N4 station that revealed low increasing rate but considerable SDDA for $T_{\rm min}$. These analyses indicated that although the northwest Tarim had greater increasing rate in $T_{\rm min}$, there were larger fluctuations in $T_{\rm min}$ of the northwest stations relative to southeast ones.

Changes in the seasonal patterns

To understand the seasonal changes, we analyzed the temperature anomalies of individual month for the past five decades (Figure 5). There were large variations between various periods except for the months from July to October. Clearly, air temperature revealed positive anomalies in all the seasons during the most recent decade, with the strongest anomaly found in March (1.54° C) and weakest one in July (0.28° C). On the other hand, there were negative anomalies in most of the months during the period of 1960–1979, with the most negative values found in the winter (-0.6 to -1.21° C). In contrast, the 1980s and 1990s had generally warm winters, with the warmest January in the 1980s and the warmest December in the 1990s. Overall, the largest decadal variability was found in early spring and late autumn.



Figure 4. Spatial distributions of linear trend of T_{min} and its SDDA during 1960–2012 in the Tarim Basin. Spatial distributions of (a) linear trend of T_{min} (°C/10a) and (b) SDDA (°C/10a) of T_{min} . doi:10.1371/journal.pone.0112231.q004

We further evaluated seasonal change by looking into the temporal variations of monthly temperature anomalies for early spring and late autumn (Figure 6). The temperature anomaly showed an increasing rate of 0.20° C/10a (P = 0.107) in March and $0.32^{\circ}C/10a$ (P = 0.004) in April. There was a similarity in temperature anomaly between March and April except during the period of 1960-1976. The temperature anomaly showed a significant increasing trend in October (0.19°C/10a, P = 0.027) and November (0.41°C/10a, P <0.001) from 1960 to 2012. Figure 6c illustrated the temperature increase in spring (i.e., the temperature difference between March and April) and decrease in autumn (i.e., the temperature difference between October and November). Overall, the temperature increase from March to April was much greater post 1970 than prior to 1970 although there was a large year-to-year variability during the most recent five years. Our data showed a general decreasing trend in the temperature change from October to November. For example, average temperature drop was 9.48°C in the 1960s but 8.30°C in the 2000s. These results implied the possibility of shortened spring and autumn with enhanced spring warming and delayed autumn cooling in the Tarim Basin since mid-1970s.

Relationships with climate indices and local environmental conditions

To explore the possible driving forcing for the temperature change in the Tarim Basin, we carried out correlation analyses, which showed that neither of PDO, SOI and DMI had significantly linear relationship with AMT, T_{min} , T_{max} or mean temperature of any season in the Tarim Basin. We also conducted non-linear relationship (data not shown). However, Table 2 showed that the correlation coefficient value was 0.264 (P = 0.059) for the relationship between the spring temperature and the SOI, suggesting that ENSO might have influence on the basin's air temperature in spring. Similarly, there was a weak negative correlation between the PDO index and the AMT in the following year (R = -0.212, P = 0.139), implying that PDO might partly be responsible for the decadal variability of air temperature in the Tarim Basin.



Figure 5. Monthly mean temperature anomalies for different decades in the Tarim Basin. doi:10.1371/journal.pone.0112231.g005

Table 3 illustrated that elevation, latitude or longitude had little influence on the warming rate of AMT. However, the warming rate of $T_{\rm min}$ showed significant relationship with elevation (R = 0.535, P = 0.018) and longitude (R = -0.528, P = 0.02), indicating greater warming in $T_{\rm min}$ at higher elevation or west stations. It seems that the significance of combined effects is about the same as that of elevation or longitude.

Discussion

Temporal and spatial variations of temperature in the Tarim Basin

In this study, a focus was placed on analyzing spatial and temporal variability of temperature change in the Tarim Basin. Our study showed that there was a large spatial variability (from



Figure 6. Time series and linear trends of monthly temperature anomalies. Trends for (a) March (black) and April (red), (b) October (black) and November (red), and (c) temperature differences between March and April (black) and between October and November (red). doi:10.1371/journal.pone.0112231.g006

| | PDO | | SOI ^b | | DMI | |
|--|---|------------------------------|------------------|-------|--------|-------|
| | ď | Р | × | μ | x | μ |
| AMT | -0.212 | 0.139 | 0.138 | 0.338 | -0.090 | 0.950 |
| T _{max} | 0.003 | 0.984 | 0.218 | 0.129 | -0.190 | 0.195 |
| T _{min} | - 0.066 | 0.647 | 0.068 | 0.640 | -0.041 | 0.784 |
| T _{spr} | -0.175 | 0.214 | 0.264 | 0.059 | -0.158 | 0.269 |
| T _{sum} | 0.110 | 0.437 | 0.023 | 0.874 | 0.156 | 0.276 |
| T _{aut} | -0.036 | 0.796 | 0.036 | 0.797 | 0.038 | 0.791 |
| T _{win} | 0.166 | 0.240 | -0.144 | 0.310 | -0.176 | 0.216 |
| ^a Temperature mea ^b Mean value over doi:10.1371/journa | ins for 1961–2012; PDO and DMI were annual m winter (e.g., from December 1960 to February 1 Lipone.0112231.t002 | eans for 1960-2011. 961). | | | | |

-0.09 to 0.43° C/10a) in the AMT warming rate during the period 1960–2012 in the Tarim Basin. The spatial variability might be a result of many factors because any single factor such as elevation could not explain the variability. The average rate (0.25°C/10a) in our study was higher than those in previous studies of the Tarim Basin (Table 3), such as 0.20° C/10a during 1960–2001 [21] and 0.22° C/10a during 1960–2007 [11]. The differences in the basin scale mean rate mainly reflected the accelerated warming (0.73°C/10a) in the most recent decade, which was consistent with the analysis by Li, Chen, Shi, Chen and Li [26] who reported a greater temperature increase (0.52°C/10a) for the period of 2000–2010 in northwest China.

An earlier study based on only six stations' data from 1958–2004 in the Tarim Basin indicated that there was a significant temperature increase in autumn and winter but not in spring and summer [15]. However, our analyses using data of 1960–2012 showed a significant warming trend in all four seasons, with the highest increasing rate in winter ($0.36^{\circ}C/10a$), and the lowest increasing rate in summer ($0.17^{\circ}C/10a$). The difference between these two studies was probably due to the significant warming in spring in the most recent decade (Figure 5), and also reflected the large spatial variability in the Tarim Basin.

Our study also showed that the warming rate was much greater in $T_{\rm min}$ (0.55°C/10a) than in $T_{\rm max}$ (0.12°C/10a) during the period 1960–2012, and there was a significant correlation between AMT and $T_{\rm min}$. These analyses indicated that air temperature increase in winter was largely responsible for the AMT increase. Overall, the seasonality had become weaker, particularly during the period of 1975–2009 in the Tarim Basin. Further analyses seemed to show an enhancement of the early spring warming and a reduction of the late autumn cooling. In addition, the increasing rate in winter (0.36°C/10a) was greater than in summer (0.17°C/10a). All these might be attributable to the weakening of the seasonality in the Tarim Basin.

Comparison between the Tarim Basin and other regions

The global mean surface air temperature had risen by $0.74\pm0.18^{\circ}$ C/100y during the twentieth century and was projected to rise by $1.8-4.0^{\circ}$ C in the twenty-first century [4,27]. For the period of 1880–2003, the linear increase of AMT over China was 0.58° C/100y, which was slightly weaker than that of the global mean [28]. However, the warming rate (0.30° C/10a) during the past two decades in China was much stronger than that of the global mean (0.19° C/10a) [29,30]. Table 4 illustrated that warming rate of the AMT was greater than 0.34° C/10a in the northwest China, but less than 0.3° C/10a in Xinjiang. Limited studies, including our study, indicated that on average, warming trend in the Tarim Basin was not as strong as that of China, which might be attributable to the smaller increase of temperature in summer.

Some studies showed an important feature associated with climate warming, i.e., the asymmetric nature over the daily cycle, with less warming observed in daily maximum temperature than in daily minimum temperature [31,32], especially in winter. A similar feature was also found over the seasonal cycle in many regions. For example, Turkes and Sumer [33] reported a significant warming of T_{min} but a weak warming and/or cooling in T_{max} in many regions of Turkey. Studies in China revealed stronger warming in T_{min} (0.32°C/10a) than in T_{max} (0.13°C/10a) [34,35]. While the increasing rate of T_{max} was comparative, the increasing rate of T_{min} was much greater in the Tarim Basin (0.55°C/10a). Similarly, based on data from 19 synoptic stations in the arid and semi-arid regions of Iran for the period of 1966–2005, Tabari [36] found that the increasing trends in the T_{min} series

Table 3. Regressions between the warming rate of temperature (AMT and T_{min}) and local variables (elevation, latitude and longitude).

| | АМТ | | T _{min} | |
|-----------------------------------|--------|-------|------------------|-------|
| Variable | R | Р | R | Р |
| Elevation | 0.149 | 0.543 | 0.535 | 0.018 |
| Latitude | -0.211 | 0.385 | 0.204 | 0.402 |
| Longitude | -0.010 | 0.968 | -0.528 | 0.020 |
| Latitude and elevation | 0.230 | 0.648 | 0.645 | 0.014 |
| Latitude and longitude | 0.210 | 0.700 | 0.612 | 0.023 |
| Longitude and elevation | 0.187 | 0.752 | 0.588 | 0.034 |
| Elevation, latitude and longitude | 0.255 | 0.791 | 0.689 | 0.019 |

doi:10.1371/journal.pone.0112231.t003

 $(0.44^{\circ}C/10a)$ were much stronger than those in the T_{max} series $(0.09^{\circ}C/10a)$. These results imply that under the global warming, arid and semi-arid regions have less extremely cold days in winter.

Impacts of local environments and large scale climate phenomena

There was evidence of warming rate increasing with the increase of latitude and/or elevation [10,37,38]. However, some studies showed a reduction in the warming rate at high elevations [39,40] or lack of clear relationship between the warming rate and elevation [41]. At the global scale, there was no simple relationship between elevation and warming rate [42]. Our analyses showed that the warming rate of AMT had little relationship with elevation (R = 0.149, P = 0.543) and latitude (R = -0.211, P = 0.385) in the Tarim Basin. However, there was a significant correlation between the warming rate of T_{min} and elevation (R = 0.535, P = 0.018), indicating greater increase of T_{min} at

higher elevation. The facts of significant relationship between AMT and $T_{\rm min}$ and great warming rate of $T_{\rm min}$ over 1960–2012 suggest that elevation may have an influence on the basin scale warming trend.

There have been limited studies that addressed the impacts of the ENSO phenomenon on the temperature in the Tarim Basin, showing inconsistent conclusions [15,17]. Using the latest dataset, our analyses indicated that ENSO events (mainly during December-February) would affect the spring temperature. Further analyses showed that there were lagged effects of both ENSO and PDO on air temperature of the Tarim Basin, i.e., ENSO on the temperature in May (R = 0.392, P = 0.004) and PDO on the temperature in July (R = 0.308, P = 0.026) of the following year.

Figure 7 illustrated that there was a large similarity in the temporal variability for the SOI, mean temperature in May and in spring, particularly post the early 1980s. In addition, the cold ENSO phases (with positive SOI, e.g., 1998–2000, and 2007)

Table 4. Increasing trends in mean air temperatures at various spatial and temporal scales.

| Region | Stations | Period | Trend | References |
|-----------------|----------|-----------|----------------|------------|
| Global | | 1901–2000 | 0.6°C/100a | [46] |
| Global | | 1906–2005 | 0.74°C/100a | [47] |
| Global | | 1956–2005 | 0.13°C/10a | [47] |
| China | 726 | 1951–2004 | 0.25°C/10a | [8] |
| China | 726 | 1905–2001 | 0.5–0.8°C/100a | [47] |
| Northern China | 486 | 1960-2000 | 0.2–0.3°C/10a | [48] |
| Southern China | 486 | 1960–2000 | <0.1°C/10a | [49] |
| Northwest China | 135 | 1960–2005 | 0.37°C/10a | [13] |
| Northwest China | 138 | 1961–2006 | 0.35°C/10a | [9] |
| Northwest China | 74 | 1960–2010 | 0.343°C/10a | [49] |
| Xinjiang | 77 | 1955–2000 | 1°C/50a | [50] |
| Xinjiang | 65 | 1961–2005 | 0.28°C/10a | [10] |
| Xinjiang | 50 | 1961–2008 | 0.30°C/10a | [51] |
| Tarim | 26 | 1960-2001 | 0.20°C/10a | [52] |
| Tarim | 23 | 1959–2006 | 0.1–0.4°C/10a | [53] |
| Tarim | 25 | 1960–2007 | 0.22°C/10a | [11] |
| Tarim | 13 | 1957–2005 | 0.19°C/10a | [12] |
| Tarim | 19 | 1960-2012 | 0.25°C/10a | This study |

doi:10.1371/journal.pone.0112231.t004

Figure 7. Time series of temperature anomalies and SOI and PDO indices in the Tarim Basin. Time series of (a) SOI and temperature anomalies in spring and May, and (b) AMT anomaly, temperature anomaly in July and PDO index. doi:10.1371/journal.pone.0112231.g007

during the most recent two decades corresponded with extremely warm temperature in May in the Tarim Basin. It seemed that there was a positive correlation between the PDO index and mean air temperature in July prior to the mid-2000s. For example, both showed an increasing trend from the early 1970s to mid-1980s, indicating a warming trend. However, for the period post late 1990s, PDO revealed mainly cold phases whereas positive anomalies (i.e., strong warming) were seen for both AMT and temperature in July.

There was evidence that the long-term climate variations in winter and summer in China might be connected to the warming trend in the sea surface temperature of the Indian Ocean [43]. Recent studies also indicated that decadal to interdecadal variability in the climate changes in Asia might be related to other climatic phenomena, such as the Arctic Oscillation [44], and Asia-Pacific Oscillation [45]. All these analyses indicated that the relationship between any of the climate indices and air temperature in the Tarim Basin might be non-linear, implying complex impacts of multi factors associated with local environmental driving and remote forcing.

Conclusions and implications

This study demonstrated a significantly increasing trend in air temperature in all four seasons during 1960–2012 in the Tarim Basin. However, there was large spatial and temporal variability in the warming rate. Temperature increase was much greater in the $T_{\rm min}$ (0.55°C/10a) than in AMT (0.25°C/10a) and $T_{\rm max}$ (0.12°C/10a), and warming rate was 0.36, 0.27, 0.23 and 0.17°C/10a in winter, autumn, spring and summer, respectively. The warming was most pronounced in the most recent decade, which might be associated with both PDO and ENSO phenomena that were in cold phases.

There was an overall weakening in the seasonality of air temperature since mid-1970s in the Tarim Basin. Apart from being less cold in winter, spring warming was another feature, which would have impacts on the hydrological cycle in the basin. Particularly, increasing temperature would lead to enhanced melting of snow and glacier in the surrounding mountains, causing extreme runoff events such as floods with a wide range of implications. Future studies are in need to better understand the climate change at various spatial scales and underlying mechanisms, and also to assess the impacts of climate change on environmental and economic aspects.

References

- Jones PD, New M, Parker DE, Martin S, Rigor IG (1999) Surface air temperature and its changes over the past 150 years. Rev Geophys 37: 173–199.
- 2. Hansen J, Ruedy R, Sato M, Lo K (2002) Global Warming Continues. Science 295: 275.
- Brown SJ, Caesar J, Ferro CAT (2008) Global changes in extreme daily temperature since 1950. J Geophys Res 113: 115–125.
 Solomon S (2007) Climate change 2007: the physical science basis: contribution
- Solomon S (2007) Climate change 2007: the physical science basis: contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge Univ Pr.
- You Q, Kang S, Aguilar E, Pepin N, Flügel WA, et al. (2011) Changes in daily climate extremes in China and their connection to the large scale atmospheric circulation during 1961–2003. Climate Dynamics 36: 2399–2417.
- Jones PD, Moberg A (2003) Hemispheric and large-scale surface air temperature variations: An extensive revision and an update to 2001. Journal of Climate 16: 206–223.
- Luterbacher J, Dietrich D, Xoplaki E, Grosjean M, Wanner H (2004) European seasonal and annual temperature variability, trends, and extremes since 1500. Science 303: 1499–1503.
- Ren G, Xu M, Chu Z, Guo J, Li Q, et al. (2005) Changes of surface air temperature in China during 1951–2004. Climatic and Environmental Research 10: 717–727.
- Chen SY, Shi YY, Guo YZ, Zheng YX (2010) Temporal and spatial variation of annual mean air temperature in arid and semiarid region in northwest China over a recent 46 year period. Journal of Arid Land 2: 87–97.
- Li Q, Chen Y, Shen Y, Li X, Xu J (2011) Spatial and temporal trends of climate change in Xinjiang, China. Journal of Geographical Sciences 21: 1007–1018.
- Xu Z, Liu Z, Fu G, Chen Y (2010) Trends of major hydroclimatic variables in the Tarim River basin during the past 50 years. Journal of Arid Environments 74: 256–267.
- Zhou H, Zhang X, Xu H, Ling H, Yu P (2011) Influences of climate change and human activities on Tarim River runoffs in China over the past half century. Environmental Earth Sciences: 1–11.
- Wang J, Fei X, Wei F (2008) Further study of temperature change in Northwest China in recent 50 years. Journal of Desert Research 4: 723–732.
- Li B, Chen Y, Li W, Chen Z, Zhang B, et al. (2013) Spatial and temporal variations of temperature and precipitation in the arid region of northwest China from 1960–2010. Fresenius Environmental Bulletin 22: 362–371.
- Chen Y, Xu C, Hao X, Weihong L, Yapeng C, et al. (2009) Fifty-year climate change and its effect on annual runoff in the Tarim River Basin, China. Quat Int 208: 53–61.
- Zhang J, Shi Y (2002) Research on Climate Change and Short-term Forecast in Xinjiang. Beijing: China Meteorological Press.
- Xu Z, Chen Y, Li J (2004) Impact of climate change on water resources in the Tarim River basin. Water Resources Management 18: 439–458.
- Chen YN, Li WH, Xu CC, Hao XM (2007) Effects of climate change on water resources in Tarim River Basin, Northwest China. Journal of Environmental Sciences-China 19: 488–493.
- Xu C, Chen Y, Li W, Chen Y, Ge H (2008) Potential impact of climate change on snow cover area in the Tarim River basin. Environmental Geology 53: 1465– 1474.
- Tao H, Gemmer M, Bai Y, Su B, Mao W (2011) Trends of streamflow in the Tarim River Basin during the past 50 years: Human impact or climate change? Journal of Hydrology 400: 1–9.
- Chen Y, Xu C, Hao X, Li W, Chen Y, et al. (2008) Fifty-year Climate Change and Its Effect on Annual Runoff in the Tarim River Basin, China. Journal of Glaciology and Geocryology 30: 921–929.
- Li Z, Yan Z (2009) Homogenized daily mean/maximum/minimum temperature series for China from 1960–2008. Atmospheric And Oceanic Science Letters 2: 237–243.
- Wu S, Yin Y, Zheng D, Yang Q (2007) Climatic trends over the Tibetan Plateau during 1971–2000. Journal of Geographical Sciences 17: 141–151.
- Király A, Jánosi IM (2005) Detrended fluctuation analysis of daily temperature records: Geographic dependence over Australia. Meteorology and Atmospheric Physics 88: 119–128.
- Talkner P, Weber RO (2000) Power spectrum and detrended fluctuation analysis: Application to daily temperatures. Physical Review E 62: 150.
- Li B, Chen Y, Shi X, Chen Z, Li W (2012) Temperature and precipitation changes in different environments in the arid region of northwest China. Theoretical and Applied Climatology: 1–8.
- Brohan P, Kennedy JJ, Harris I, Tett SFB, Jones PD (2006) Uncertainty estimates in regional and global observed temperature changes: A new data set from 1850. Journal of geophysical research 111: 106–126.

Author Contributions

Conceived and designed the experiments: XJW XRW. Performed the experiments: DMP. Analyzed the data: DMP PXC. Contributed reagents/ materials/analysis tools: CYZ FQJ. Wrote the paper: DMP XJW.

- Wang S, Gong D (2000) Enhancement of the warming trend in China. Geophysical Research Letters 27: 2581–2584.
- Wang S, Zhu J, Cai J (2004) Interdecadal variability of temperature and precipitation in China since 1880. Advances in Atmospheric Sciences 21: 307– 313.
- Hansen J, Sato M, Ruedy R, Lo K, Lea DW, et al. (2006) Global temperature change. Proceedings of the National Academy of Sciences 103: 14288–14293.
- Karl TR, Jones PD, Knight RW, Kukla G, Plummer N, et al. (1993) Asymmetric trends of daily maximum and minimum temperature. Papers in Natural Resources 74: 1007–1023.
- Liu X, Cheng Z, Yan L, Yin ZY (2009) Elevation dependency of recent and future minimum surface air temperature trends in the Tibetan Plateau and its surroundings. Global and Planetary Change 68: 164–174.
- Türkeş M, Sümer U (2004) Spatial and temporal patterns of trends and variability in diurnal temperature ranges of Turkey. Theoretical and Applied Climatology 77: 195–227.
- Liu B, Xu M, Henderson M, Qi Y, Li Y (2004) Taking China's Temperature: Daily Range, Warming Trends, and Regional Variations, 1955–2000. Journal of climate 17: 4453–4462.
- Wei F (2008) Probability Distribution of Minimum Temperature in Winter Half Years in China. Advances in Climate Change Research 4: 8–11.
- Tabari H, Hosseinzadeh Talaee P (2011) Analysis of trends in temperature data in arid and semi-arid regions of Iran. Global and Planetary Change 79: 1–10.
- Holmes JA, Cook ER, Yang B (2009) Climate change over the past 2000 years in Western China. Quat Int 194: 91–107.
- Wang G, Shen Y, Zhang J, Wang S, Mao W (2010) The effects of human activities on oasis climate and hydrologic environment in the Aksu River Basin, Xinjiang, China. Environmental Earth Sciences 59: 1759–1769.
- Pepin N, Losleben M (2002) Climate change in the Colorado Rocky Mountains: free air versus surface temperature trends. International Journal of Climatology 22: 311–329.
- Vuille M, Bradley RS (2000) Mean annual temperature trends and their vertical structure in the tropical Andes. Geophys Res Lett 27: 3885–3888.
- Pepin N, Lundquist J (2008) Temperature trends at high elevations: patterns across the globe. Geophysical Research Letters 35: L14701.
- Pepin N, Seidel DJ (2005) A global comparison of surface and free-air temperatures at high elevations. Journal of Geophysical Research 110: D03104.
- Hu ZZ, Yang S, Wu RG (2003) Long-term climate variations in China and global warming signals. Journal of Geophysical Research-Atmospheres 108.
- 44. Woo SH, Kim BM, Jeong JH, Kim SJ, Lim GH (2012) Decadal changes in surface air temperature variability and cold surge characteristics over northeast Asia and their relation with the Arctic Oscillation for the past three decades (1979–2011). Journal of Geophysical Research-Atmospheres 117.
- Zhao P, Yang S, Wang HJ, Zhang Q (2011) Interdecadal Relationships between the Asian-Pacific Oscillation and Summer Climate Anomalies over Asia, North Pacific, and North America during a Recent 100 Years. Journal of Climate 24: 4793–4799.
- Xu J, Li W, Ji M, Lu F, Dong S (2010) A comprehensive approach to characterization of the nonlinearity of runoff in the headwaters of the Tarim River, western China. Hydrological processes 24: 136–146.
- Ding Y, Ren G, Zhao Z, Xu Y, Luo Y, et al. (2007) Detection, causes and projection of climate change over China: an overview of recent progress. Advances in Atmospheric Sciences 24: 954–971.
- Qian W, Qin A (2006) Spatial-temporal characteristics of temperature variation in China. Meteorology and Atmospheric Physics 93: 1–16.
- Li B, Chen Y, Shi X (2012) Why does the temperature rise faster in the arid region of northwest China? Journal of Geophysical Research: Atmospheres (1984–2012) 117.
- Hao X, Chen Y, Xu C, Li W (2008) Impacts of climate change and human activities on the surface runoff in the Tarim River basin over the last fifty years. Water Resources Management 22: 1159–1171.
- Zhang Y, Wei W, Jiang F, Liu M, Wang W, et al. (2012) Brief communication "Assessment of change in temperature and precipitation over Xinjiang, China". Natural Hazards and Earth System Sciences 12: 1327–1331.
- Liu Z, Xu Z (2007) Spatio-temporal distribution of hydrometeorological variables and their main impact factors in the Tarim river basin. Journal of China Hydrology 5: 020.
- Xu J, Chen Y, Li W, Ji M, Dong S, et al. (2009) Wavelet analysis and nonparametric test for climate change in Tarim River Basin of Xinjiang during 1959–2006. Chinese Geographical Science 19: 306–313.