

Jishao Jiang^{1,2}
 Rui Wang³
 Zhiqi Wang⁴
 Shengli Guo^{1,3,4}
 Xiaotang Ju^{5*}

Research Article

Nitrous Oxide and Methane Emissions in Spring Maize Field in the Semi-Arid Regions of Loess Plateau

¹Institute of Soil and Water Conservation, Chinese Academy of Sciences and Ministry of Water Resources, Yangling, Shaanxi, P. R. China

²Key Laboratory for Yellow River and Huai River Water Environmental and Pollution Control, Ministry of Education, Henan Key Laboratory for Environmental Pollution Control, School of Environment, Henan Normal University, Xinxiang, Henan, P. R. China

³College of Resources and Environment, Northwest A&F University, Yangling, P.R. China

⁴State Key Laboratory of Soil Erosion and Dryland Farming on the Loess Plateau, Institute of Soil and Water Conservation, Northwest A&F University, Yangling, P. R. China

⁵College of Resources and Environmental Sciences, China Agricultural University, Beijing, P. R. China

A 2-year field study was conducted to measure nitrous oxide (N₂O) and methane (CH₄) in a rain-fed spring maize cropland in the Loess Plateau, P. R. China, and to determine the effects of optimized nitrogen (N) fertilization practices on urea-derived N₂O emission factor (EF), grain yield, net greenhouse gas (NGHG) emission, and net greenhouse gas intensity (NGHGI). Five treatments were considered, including control (CK), conventional N fertilization (Con), optimal N fertilization (Opt), optimal N fertilization plus nitrification inhibitor (Opt + DCD), and optimal N fertilization with slow release urea (Opt + SR). Soil acted as a small sink for atmospheric CH₄. Nitrogen fertilization and heavy rainfall events (>40 mm) were the main factors controlling N₂O emissions. The annual mean EF ranged from 0.12 to 0.55%. Compared to conventional N fertilizer, nitrification inhibitor decreased the annual cumulative N₂O, NGHG, and NGHGI emissions by 45, 52, and 48%, respectively, without decreasing grain yield. In conclusion, nitrification inhibitor addition was the most effective practice to reduce N₂O emissions in the rain-fed regions of Loess Plateau.

Keywords: Nitrification inhibition; Nitrogen practice; Nitrous oxide; Slow release urea; Spring maize

Received: April 19, 2015; *revised:* April 5, 2016; *accepted:* October 19, 2016

DOI: 10.1002/clean.201500271

1 Introduction

Agriculture constitutes an important source of nitrous oxide (N₂O) and methane (CH₄) [1], accounting for approximately 60 and 50% of global N₂O and CH₄ emissions, respectively [2]. Global annual N₂O emissions from nitrogen (N) fertilizers and manures, crop residues, and other agricultural sources are estimated to be up to 4.1 Tg N₂O–N y⁻¹ (range: 3.8–6.8) [3]. Thus, there is an urgent need to reduce N₂O and CH₄ emissions from croplands to mitigate global greenhouse gas (GHG) emissions.

N₂O emissions from cropland increase with an increase in N fertilizer rate [4–6]. Many N fertilization practices, such as balanced N fertilization [7, 8], nitrification inhibitor (NI) [8, 9], and slow release (SR) urea [10, 11], have been proposed to reduce GHG emissions from

N-fertilized croplands. However, it is important to note that the effectiveness of these N fertilization practices in reducing N₂O emissions differs greatly, depending on the environmental conditions and farming systems. For instance, some studies showed that NI reduced N₂O emissions by 50–82% in comparison with conventional urea [12, 13]; whereas other studies showed that NI had a limited impact on N₂O emissions [14, 15]. Similarly, SR was found to be able to significantly reduce N₂O emissions from grassland in UK [16], but increase N₂O emissions in the wheat and maize croplands in eastern China [11]. These conflicting results indicate that the mechanism underlying the effects of NI and SR on N₂O emissions remains poorly understood. In addition, upland soils act as a weak sink for atmospheric CH₄ [17], but N fertilization has a variable effect on CH₄ emissions [18]. Some studies reported that N application inhibited atmospheric CH₄ uptake by soils [8, 17], whereas other studies concluded that fertilizer N rates had no significant effect on CH₄ uptake [19]. Thus, more field experiments are required to better understand the effect of different N fertilization practices on N₂O and CH₄ emissions.

GHG emissions are strongly controlled not only by temperature, but also by soil moisture and soil mineral N [20–23]. The content and transformation of soil mineral N could directly influence

Correspondence: Shengli Guo, Institute of Soil and Water Conservation, Chinese Academy of Sciences and Ministry of Water Resources, Yangling, Shaanxi 712100, P. R. China
 E-mail: slguo@ms.iswc.ac.cn

Abbreviations: ANOVA, one-way analysis of variance; CK, control; Con, conventional; CV, coefficient of variation; DCD, dicyandiamide; ECD, electron capture detector; EF, emission factor; FID, flame ionization detector; GHG, greenhouse gas; IPCC, Intergovernmental Panel on Climate Change; LSD, least significant difference; NGHG, net greenhouse gas emission; NGHGI, NGHG intensity; NI, nitrification inhibitor; Opt, optimal; SR, slow release; TN, total nitrogen; WFPS, soil water-filled pore space.

^{*}Additional correspondence: Prof. Xiaotang Ju, College of Resources and Environmental Sciences, China Agricultural University, Beijing, P. R. China. E-mail: juxt@cau.edu.cn

nitrification and denitrification [5]. In the arid and semi-arid regions, rainfall caused the fluctuation of soil moisture, thus, affecting the microbial activity related to N_2O and CH_4 emissions and the transformation of soil mineral N, and consequently the N_2O and CH_4 emissions [11, 21]. The emission factor (EF) was used to estimate N_2O emissions on the basis of the amount of N fertilizer used in a country or geographic region, usually using the default factor (1 or 1.25% derived from Intergovernmental Panel on Climate Change (IPCC) [24] and a meta-analysis of EF in global cropland [25], respectively). However, it is noted that arid and semi-arid regions are not included in the above meta-analysis. In addition, previous empirical studies have shown that EF is not constant, but varies from 0.1 to 73.7% in different crops, countries, and regions [26]. Thus, an accurate EF is needed for accurate estimation of N_2O emissions in arid and semi-arid regions.

In the Loess Plateau, P. R. China, the soil fertility in general and the soil N level in particular are very low, with a total nitrogen (TN) content of 0.06–0.08% [27, 28]. Therefore, winter wheat and spring maize, the two main crops in this region, are heavily fertilized with chemical N fertilizer (200 and 230 kg N ha⁻¹, respectively) during the past decades to obtain a high yield [29–31]. Rainfall is the sole source of soil water supply. In this region, the annual precipitation for the period 1995–2004 ranged from a minimum of 332 mm in 1995 to a maximum of 919 mm in 2004 with a co-efficient of variation (CV) of 22%, and the precipitation from July to September ranged from 147 to 609 mm with a CV of 37% [32]. As the effect of N fertilization depends critically on water availability in the arid and semi-arid regions [33, 34], N fertilizer tends to be of low efficiency in the Loess Plateau, resulting in potential environmental risks such as GHG emissions [35, 36] and subsoil nitrate accumulation and leaching [37, 38]. However, there have been few studies investigating EF and the main factors controlling N_2O and CH_4 emissions in rain-fed farming regions.

N_2O and CH_4 emissions from a rain-fed spring maize cropland in the semi-arid Loess Plateau were measured in this 2-year field study. The objectives were to (i) quantify N_2O and CH_4 emissions after various N fertilization practices; (ii) identify EFs and the main factors controlling N_2O and CH_4 emissions; and (iii) identify the most effective N fertilization practice in the semi-arid Loess Plateau, P. R. China.

2 Materials and methods

2.1 Site description

A field experiment was carried out at State Key Agro-Ecological Experimental Station in the Loess Plateau (35°12'N, 107°40'E; 1220 m asl) in Changwu County, Shaanxi Province, P. R. China. The study area is a typical rain-fed farming region and has a semi-arid continental monsoon climate. The mean annual rainfall from 1984 to 2014 was 560 mm, about 60% of which falls between June and September. The open pan evaporation is 1440 mm. The mean annual temperature is 9.4°C. The daily air temperature and precipitation during the study period from January 2013 to December 2014 are shown in Fig. 1.

The soil at the study site is loam (Cumulic Haplustoll; USDA Soil Taxonomy System) developed from loess deposits, and contains 8% sand, 70% silt, and 22% clay [39]. Soils in the top 20 cm are composed of $CaCO_3$ 10.5%, organic C 6.5 g kg⁻¹, and TN 0.80 g kg⁻¹, with a field water holding capacity of 0.29 cm³ cm⁻³,

a pH of 8.4 (soil: H₂O suspension, 1:1), and a bulk density of 1.3 Mg m⁻³.

2.2 Experimental design and crop management

A high-yield spring maize (*Zea mays* L.) hybrid (Pioneer, 335), a representative crop type in this region, was chosen for this field study. There were five treatments with three replicates each in a completely randomized design, giving a total of 15 plots. All plots were 5.5 × 18 m with a spacing of 0.5 m between adjacent plots and 1 m between adjacent blocks. The treatments were as follows:

- (1) Control (CK): No fertilizer N applied.
- (2) Conventional N fertilizer (Con): Urea (N 46%) applied at a rate of 200 kg N ha⁻¹ y⁻¹, which is the general practice in this region.
- (3) Optimum N fertilizer (Opt): Urea applied at a rate of 160 kg N ha⁻¹ y⁻¹ based on the recommendation of local extension service.
- (4) Optimum N fertilizer plus NI (Opt + DCD): Urea and solid dicyandiamide (DCD, chemically pure, Sinopharm Chemical Reagents) applied at a rate of 10% N, and broadcast onto the soil surface before plough.
- (5) Optimum N fertilizer with slow release urea (Opt + SR): SR is coated urea (urea formaldehyde) containing 26% N.

Phosphorus (superphosphate) and potassium (potassium sulfate) fertilizers were applied at a rate of 117 kg P₂O₅ ha⁻¹ and 37.5 kg K₂O ha⁻¹ before seeding. All fertilizers were mixed into the top 20 cm soil as basal fertilizer by ploughing. Soil was covered with plastic film (750 mm wide by 0.008 mm thick) in order to reduce soil evaporation and increase soil temperature, and crop growth. The plant density was 57 000 plants ha⁻¹ at a distance of 30 cm in rows and 60 cm between rows.

Maize straw was removed from the plots after harvest, and weeds were removed manually during the growing season. No irrigation was applied during the growing season, and thus, soil water supply was completely provided by natural rainfall. Fallow (from September to May of the next year) and maize (from May to September) were rotated at the study site. Maize was planted on April 24 and harvested on September 9 in 2013, and planted on April 30 and harvested on September 15 in 2014.

2.3 Gas sampling and measurements

N_2O and CH_4 emissions were measured using the static chamber method [40]. Two types of static chambers were used in this study, each consisting of base frame and a removable upper chamber made of stainless steel. The base frame and chamber were 60 × 50 × 20 cm (length × width × height) and 60 × 50 × 50 cm for type I chamber; and 60 × 30 × 20 cm and 60 × 30 × 20 cm (length × width × height) for type II chamber, respectively. Prior to planting, the base frames were inserted 20 cm into the soil (half mulching soil and half bare soil), and then remained in place throughout the measurement period except for tillage, which was performed once a year. A visual description is shown in Fig. 2. Type I chamber was used to measure N_2O emissions when maize height was <50 cm, with two maize plants in each chamber; while type II chamber was used when maize height was >50 cm. Type II chamber was separated vertically into two parts with a hole (11 cm in diameter) at the top center of the chamber, and only one maize plant was placed in the chamber [41].




Figure 1. Air and soil temperature at 0–20 cm depths (a) and precipitation and soil WFPS at 0–20 cm depths (b). (FS, fallow season; MS, maize growing season).

This allowed the cornstalks to pass through the chamber and as a result, to only cover the maize root. The gap between the chamber and the cornstalk was sealed using a 1.2 μm -thick preservative film made of polyvinylidene chloride when the chamber was closed. All chambers were sealed with rubber and covered with an insulating layer to minimize chamber effects on air temperature to $<3^\circ\text{C}$ in the headspace during gas sampling, and two opposing ventilators were installed inside the chamber to ensure complete mixing of air.

Gas samples were collected using 50 mL plastic injectors through a three-way stopcock and a Teflon tube connected to the chamber at 0, 10, 20, 30, 40 min after the chambers were closed. Under normal circumstances, N_2O and CH_4 emissions were measured once a week during the growing season of maize, once a month when the soil was frozen (from November to March of next year), and every 10 days for the fallow season. However, daily measurements were carried out for about 10 days after fertilizer application and 3 days after heavy rainfall events ($>20\text{ mm}$), respectively. All measurements were conducted between 08:00 a.m. and 11:00 a.m. local time, and there were a total of 57 and 40 measurements in 2013 and 2014, respectively.

Gas samples were analyzed for N_2O and CH_4 within 24 h of sampling using gas chromatography (Agilent GC6820, Agilent, USA) equipped with an electron capture detector (ECD) and a flame ionization detector (FID). Temperatures in the ECD and column oven were 300 and 60°C , respectively. Pure N_2 (99.999%) was used as the carrier gas and 10% CO_2 in pure N_2 as the buffer gas for ECD, respectively. The detection limit was $2\ \mu\text{g N m}^{-2}\text{ h}^{-1}$ for N_2O and $3.4\ \mu\text{g C m}^{-2}\text{ h}^{-1}$ for CH_4 when the chamber height was 50 cm. The gas samples were calibrated using compressed air ($333\ \mu\text{L m}^{-3}\text{ N}_2\text{O}$ and $197\ \mu\text{L m}^{-3}\text{ CH}_4$) during each measurement cycle.

N_2O and CH_4 emission rates were calculated from the linear increase in the concentration in the chamber during the sampling

period. As all measurements of CH_4 emissions were negative, CH_4 uptakes (positive) were used for convenience in this study. The cumulative N_2O emissions and CH_4 uptakes were estimated using linear interpolation between every two adjacent intervals of the measurements (Eqs. (1) and (2)) [37].

$$N_{\text{cum}} = \sum_{i=\text{first}}^{i=\text{last}-1} \left(\left(\frac{D_i - D_{i+1}}{2} \right) \times (N_{i+1} - N_i - 1) + D_i \right) + D_{\text{last}} \quad (1)$$

$$D_i = F \times 24 \times 10^{-5} \quad (2)$$

where N_{cum} ($\text{kg ha}^{-1}\text{ y}^{-1}$) is the annual N_2O cumulative emissions or CH_4 cumulative uptakes; D_i (kg ha^{-1} per day) is the daily N_2O cumulative emissions or CH_4 cumulative uptakes; first and last are the first and last measurements of N_2O emissions or CH_4




Figure 2. Schematic diagram of the installation of base frames. Black dots denote the maize plants. Shadow denote the plastic film. Types I and II denote the two types of chambers used in this study.

uptakes; $N_{i+1}-N_i-1$ is the interval between two adjacent measurements; F ($\mu\text{g m}^{-2} \text{h}^{-1}$) is the N_2O emission rate or CH_4 uptake rate; 24 and 10^{-5} are the conversion co-efficients.

2.4 Soil temperature, soil moisture, $\text{NO}_3\text{-N}$, and $\text{NH}_4\text{-N}$

Air temperature inside the chamber and soil temperature at a 10 cm depth were measured using a portable digital thermometer (JM624, Tianjin Jinming Instruments, P. R. China) immediately after the first and last sampling, and the mean of the two measurements was used as the temperature on the sampling day. Soil samples at a 20 cm depth were collected once every 2 days for 10 days following N fertilization and within 4 days following heavy rainfalls. Over the remaining period, soils were sampled once when gases were sampled. Gravimetric soil water content w/w was measured after drying the soil in an oven at 105°C for 24 h, and soil water-filled pore space (WFPS) was calculated using Eq. (3). To determine $\text{NH}_4\text{-N}$ and $\text{NO}_3\text{-N}$ content, fresh samples (24 g) were extracted using 100 mL of 1 mol L^{-1} KCl solution, and then the extracts were analyzed using continuous flow analysis (TRAACS2000, Bran and Luebbe, Norderstedt, Germany).

$$\text{WEPS} = \frac{\text{Soil water content (\%)} \times \text{Soil bulk density}}{1 - \frac{\text{Soil bulk density}}{2.65}} \times 100\% \quad (3)$$

2.5 Grain yield

Maize was manually harvested from an area of 16 m^2 in each plot. The samples were dried at 65°C to a constant weight to determine the aboveground biomass. The grain yield was expressed at 15.5% moisture [42].

2.6 Net greenhouse gas emission (NGHG), net greenhouse gas emission intensity (NGHGI), and EF

NGHG emissions were calculated in terms of N_2O plus CH_4 fluxes in CO_2 equivalents ($\text{CO}_2\text{-eq}$) to evaluate the effects of different N fertilization practices on N_2O emissions and CH_4 uptakes. The global warming potentials of $1 \text{ kg N}_2\text{O}$ and CH_4 being equivalent to 298 and

25 kg CO_2 at the 100-year time horizon, respectively [41], were applied to measure NGHG ($\text{kg CO}_2\text{-eq ha}^{-1}$) in $\text{CO}_2\text{-eq}$. NGHGI ($\text{kg CO}_2\text{-eq Mg}^{-1}$ grain) was calculated by dividing NGHG by crop yield and expressed as the magnitude of NGHG to produce the same crop yield. EF was determined using Eq. (4).

$$\text{EF} = \sum_{i=\text{first}}^{i=\text{last}-1} \left(\left(\frac{D_i - D_{i+1}}{2} \right) \times (N_{i+1} - N_i - 1) + D_i \right) + D_{\text{last}}$$

$$\text{EF} = \frac{N_{\text{tr-cum}} - N_{\text{ck-cum}}}{N_{\text{rate}}} \times 100\% \quad (4)$$

where $N_{\text{tr-cum}}$ ($\text{kg N}_2\text{O-N ha}^{-1} \text{y}^{-1}$) is the annual N_2O cumulative emissions in the four N treatments, $N_{\text{ck-cum}}$ ($\text{kg N}_2\text{O-N ha}^{-1} \text{y}^{-1}$) is the annual N_2O cumulative emissions in CK treatment, and N_{rate} ($\text{kg N ha}^{-1} \text{y}^{-1}$) is the N fertilization rate in the four N treatments.

2.7 Data analysis

All data in this study are presented as mean \pm SD. In each year, the differences in grain yield, seasonal, and annual N_2O and CH_4 cumulative emissions among the five treatments were analyzed using one-way analysis of variance (ANOVA) followed by least significant difference (LSD) post-hoc test. A p -value of <0.05 was considered significant. Pearson correlation analysis was performed to determine the relationships between N_2O emissions and soil variables. All statistical analyses were performed using SPSS 17.0 for Windows (SPSS Statistics, Chicago, IL, USA).

3 Results

3.1 Grain yield

Nitrogen fertilization significantly enhanced maize grain yield in both 2013 and 2014 ($p < 0.05$; Tab. 1). The maize yield for Opt, Opt + DCD, and Opt + SR treatments ranged from 9.61 to 10.46 Mg ha^{-1} in 2013 with an increase of 31.6–43.3%; and from 11.41 to 11.92 Mg ha^{-1} in 2014 with an increase of 185–198% in comparison to CK treatment, respectively. However, there was no significant difference in the grain yield between N optimized (Opt, Opt + DCD, and Opt + SR) treatments and Con treatment (11.3 Mg ha^{-1}) in both 2013 and 2014, although the N fertilization rate of the former was 20–25% lower than that of the latter.

Table 1. Yield, NGHG, and NGHGI for different N fertilization practices in 2013 and 2014

Treatment	2013			2014		
	Yield (Mg ha^{-1})	NGHG ($\text{kg CO}_2\text{-eq ha}^{-1}$)	NGHGI ($\text{kg CO}_2\text{-eq Mg}^{-1}$ grain)	Yield (Mg ha^{-1})	NGHG ($\text{kg CO}_2\text{-eq ha}^{-1}$)	NGHGI ($\text{kg CO}_2\text{-eq Mg}^{-1}$ grain)
N0	7.30 ± 0.94^b	301 ± 15^d	42 ± 4^{bc}	3.99 ± 1.41^b	240 ± 11^d	65 ± 22^a
Con	10.30 ± 0.44^a	807 ± 53^a	78 ± 4^a	12.23 ± 0.66^a	769 ± 74^a	63 ± 9^a
Opt	10.46 ± 0.33^a	555 ± 72^b	53 ± 8^b	11.63 ± 2.69^a	517 ± 53^b	47 ± 18^{ab}
Opt + DCD	9.61 ± 0.84^a	383 ± 59^{cd}	40 ± 9^c	11.92 ± 0.94^a	304 ± 56^c	26 ± 6^b
Opt + SR	10.10 ± 0.75^a	444 ± 51^c	44 ± 7^{bc}	11.41 ± 0.54^a	438 ± 51^b	38 ± 4^b

$\text{NGHG} (\text{kg CO}_2\text{-eq ha}^{-1}) = \text{N}_2\text{O-N} \times 44/28 \times 298 + \text{CH}_4\text{-C} \times 16/12 \times 25$.

$\text{NGHGI} (\text{kg CO}_2\text{-eq Mg}^{-1} \text{ grain}) = \text{NGHG}/\text{yield}$.

Values are expressed as mean \pm standard deviation ($n = 3$). The values within the columns followed by different letters are significantly different at $p < 0.05$.




Figure 3. Nitrous oxide emissions (N_2O) from fields with different nitrogen fertilization practices across fallow and maize growing season in 2013 and 2014. (FS, fallow season; MS, maize growing season).

3.2 N_2O emissions

In both years, the N_2O emission rates in all treatments increased rapidly to a peak within 2 days after N fertilization, maintained at a high level for about 10 days, and then decreased rapidly to a lower level of $0\text{--}20 \mu\text{g } N_2O\text{--}N \text{ m}^{-2} \text{ h}^{-1}$ (Fig. 3). However, peaks could also be observed after high rainfalls ($>40 \text{ mm}$). During the 2-year study period, the mean N_2O emission rate was 28.8 , 17.3 , and $21.4 \mu\text{g } N_2O\text{--}N \text{ m}^{-2} \text{ h}^{-1}$ for Opt, Opt + DCD, and Opt + SR treatments, with a decrease of 29.5 , 57.7 , and 47.7% compared with Con treatment ($40.9 \mu\text{g } N_2O\text{--}N \text{ m}^{-2} \text{ h}^{-1}$), respectively. In both years, the cumulative N_2O emissions within 10 days after N fertilization accounted for about 13% of the total annual N_2O emissions in CK and Opt + DCD treatments, and about 26% in other treatments, respectively. In 2013, the largest daily N_2O emission was observed on the second day of N fertilization, which was 26.5 , 160.7 , 143.1 , 52.3 , and $77.7 \mu\text{g } N_2O\text{--}N \text{ m}^{-2} \text{ h}^{-1}$ in CK, Con, Opt, Opt + DCD, and Opt + SR

treatments, respectively. A similar phenomenon was also observed in 2014. N_2O emissions had a strong response to rainfall, especially when the rainfall was $>40 \text{ mm}$. For example, N_2O emission peaks were observed after heavy rainfall on July 22, 2013 (120 mm) (31.4 , 77.7 , 58.5 , 33.0 , and $26.9 \mu\text{g } N_2O\text{--}N \text{ m}^{-2} \text{ h}^{-1}$ in CK, Con, Opt, Opt + DCD, and Opt + SR, respectively). A similar phenomenon was also observed on August 6 (44 mm) and September 30 (40.6 mm) in 2014. The cumulative N_2O emissions induced by rainfall accounted for 6.4 and 12.5% of the total annual N_2O emissions in 2013 and 2014, respectively.

The annual N_2O emissions in Con, Opt, Opt + DCD, and Opt + SR treatments ($0.91\text{--}1.88 \text{ kg } N_2O\text{--}N \text{ ha}^{-1} \text{ y}^{-1}$) were significantly higher than that of CK treatment ($0.80 \text{ kg } N_2O\text{--}N \text{ ha}^{-1} \text{ y}^{-1}$) ($p < 0.05$), indicating that N fertilization could significantly increase annual N_2O emissions (Tab. 2). However, the annual N_2O emission in Opt, Opt + DCD, and Opt + SR treatments was decreased by 27.4 , 45.2 , and 39.6% in 2013, and by 27.7 , 51.6 , and

Table 2. Cumulative N₂O emission (kg N₂O–N ha⁻¹) and CH₄ uptake (kg CH₄–C ha⁻¹) for different N fertilization practices in 2013 and 2014

Treatment	2013			2014		
	MS	FS	MS + FS	MS	FS	MS + FS
N₂O						
NO	0.52 ± 0.01 ^d	0.35 ± 0.01 ^b	0.87 ± 0.03 ^d	0.39 ± 0.02 ^c	0.33 ± 0.01 ^b	0.72 ± 0.03 ^c
Con	1.42 ± 0.12 ^a	0.54 ± 0.05 ^a	1.97 ± 0.09 ^a	1.28 ± 0.15 ^a	0.60 ± 0.06 ^a	1.88 ± 0.14 ^a
Opt	1.01 ± 0.10 ^b	0.42 ± 0.05 ^b	1.43 ± 0.14 ^b	0.90 ± 0.06 ^b	0.46 ± 0.06 ^b	1.36 ± 0.12 ^b
Opt + DCD	0.68 ± 0.06 ^c	0.40 ± 0.05 ^b	1.08 ± 0.11 ^c	0.50 ± 0.07 ^c	0.41 ± 0.03 ^b	0.91 ± 0.10 ^c
Opt + SR	0.79 ± 0.11 ^c	0.39 ± 0.01 ^b	1.19 ± 0.11 ^c	0.77 ± 0.13 ^b	0.43 ± 0.01 ^b	1.20 ± 0.13 ^b
CH₄						
NO	1.88 ± 0.12 ^a	1.31 ± 0.15 ^a	3.19 ± 0.26 ^a	1.65 ± 0.02 ^a	1.32 ± 0.08 ^a	2.98 ± 0.09 ^a
Con	1.84 ± 0.01 ^a	1.61 ± 0.27 ^a	3.45 ± 0.26 ^a	1.57 ± 0.06 ^a	1.78 ± 0.31 ^a	3.35 ± 0.28 ^a
Opt	1.80 ± 0.05 ^a	1.66 ± 0.14 ^a	3.46 ± 0.17 ^a	1.86 ± 0.07 ^a	1.74 ± 0.27 ^a	3.60 ± 0.23 ^a
Opt + DCD	1.93 ± 0.05 ^a	1.74 ± 0.32 ^a	3.67 ± 0.28 ^a	1.91 ± 0.05 ^a	1.73 ± 0.32 ^a	3.64 ± 0.37 ^a
Opt + SR	1.62 ± 0.10 ^a	1.73 ± 0.05 ^a	3.35 ± 0.11 ^a	1.83 ± 0.09 ^a	1.89 ± 0.13 ^a	3.72 ± 0.09 ^a

FS, fallow season; MS, maize growing season.

Values are expressed as mean ± standard deviation (*n* = 3). The values within the columns followed by different letters are significantly different at *p* < 0.05 in N₂O or CH₄.

36.2% in 2014 compared with Con treatment (1.88 kg N₂O–N ha⁻¹ in 2013 and 1.97 kg N₂O–N ha⁻¹ in 2014), respectively. The cumulative N₂O emission during the fallow season accounted for 27.4–40.2% of the total annual N₂O emissions in 2013 and 31.9–45.8% in 2014.

EF values ranged from 0.12 to 0.55%, and were higher than that of growth season (0.07–0.45%). The lowest value was observed in Opt + DCD treatment, followed by Opt + SR, Opt, and Con treatments (Tab. 3).

3.3 CH₄ uptakes

Soils acted as a small sink for atmospheric CH₄ (Fig. 4). There was no significant difference in daily or annual mean CH₄ uptakes among the five treatments (*p* > 0.05). The daily CH₄ uptake ranged from 0.15 to 127 μg CH₄–C m⁻² h⁻¹ with an average of 45 μg CH₄–C m⁻² h⁻¹, and the annual CH₄ uptake ranged from 3.0 to 3.7 kg CH₄–C ha⁻¹ with an average of 3.4 kg CH₄–C ha⁻¹ (Tab. 2). CH₄ uptakes increased from April to July and then decreased slowly in winter.

3.4 Soil temperature, WFPS, NO₃–N, and NH₄–N

Soil temperature at a 10 cm depth (range: –0.5 to 24.7°C; mean: 17.7°C) varied with daily mean air temperature (Fig. 1a). In the top 20 cm soils, soil WFPS markedly increased after heavy rainfall events and then decreased rapidly due to soil texture, high evaporation,

and crop uptake (Fig. 1b). It ranged from 17.0 to 70.0% with an average of 48.8%, and was <60.0% for most of the study period.

The NO₃–N content showed a high response to N fertilization and heavy rainfalls (>40 mm) (Fig. 5). The average soil NO₃–N content was 22, 106, 100, 71, and 61 mg kg⁻¹ for CK, Con, Opt, Opt + DCD, and Opt + SR treatments, respectively. The NO₃–N content increased markedly after N fertilization, and maintained at high level for a relatively long period. Application of NI and SR led to a lower NO₃–N content than Opt treatment. Furthermore, the NO₃–N content increased rapidly after heavy rainfalls. For example, the NO₃–N content in all treatments except CK treatment reached a peak after the heavy rainfall (120 mm) on 22 July, 2013 (217, 276, 37, and 83 mg kg⁻¹ for Con, Opt, Opt + DCD, and Opt + SR treatments, respectively). The same phenomenon was also observed after a heavy rainfall (44 mm) on August 6, 2014. In contrast, the NH₄–N content was relatively low (0.35 to 58.0 mg kg⁻¹) (Fig. 6). It increased rapidly to a peak within 2 days after fertilization, and then decreased rapidly to a lower level. The mean NH₄–N content was 5, 13, 10, 12, and 9 mg kg⁻¹ for CK, Con, Opt, Opt + DCD, and Opt + SR treatments, respectively. Application of NI and SR led to higher soil NH₄–N content than Opt treatment.

3.5 NGHG and NGHGI

Compared with Con treatment, NGHG was decreased by 31.2, 52.5, and 45.0% in Opt, Opt + DCD, and Opt + SR treatments in 2013, and

Table 3. Direct N₂O emission factors (%) for different N fertilization practices in 2013 and 2014

Treatments	2013		2014		Mean	
	MS	MS + FS	MS	MS + FS	MS	MS + FS
Con	0.45 ± 0.05 ^a	0.55 ± 0.05 ^a	0.44 ± 0.07 ^a	0.58 ± 0.07 ^a	0.45	0.57
Opt	0.31 ± 0.05 ^b	0.35 ± 0.08 ^b	0.32 ± 0.03 ^b	0.40 ± 0.07 ^b	0.31	0.38
Opt + DCD	0.10 ± 0.04 ^c	0.13 ± 0.07 ^c	0.07 ± 0.04 ^c	0.12 ± 0.06 ^c	0.08	0.12
Opt + SR	0.17 ± 0.07 ^c	0.20 ± 0.07 ^c	0.24 ± 0.07 ^b	0.30 ± 0.07 ^b	0.20	0.25

FS, fallow season; MS, maize growing season.

Values are expressed as mean ± standard deviation (*n* = 3). The values within the columns followed by different letters are significantly different at *p* < 0.05.




Figure 4. Methane (CH₄) uptake from fields with different nitrogen fertilization practices across fallow and maize growing season in 2013 and 2014. (FS, fallow season; MS, maize growing season).

by 32.8, 60.5, and 43.0% in 2014 ($p < 0.05$) (Tab. 1), respectively; and NGHGI was decreased by 32.1, 48.7, and 43.6% in 2013, and by 25.4, 58.7, and 39.7% in 2014, respectively. The highest decrease occurred in Opt + DCD treatment, followed by Opt + SR treatment.

4 Discussion

4.1 N₂O emissions from croplands in semi-arid regions

The annual cumulative N₂O emissions after N fertilization practices ranged from 0.72 to 1.97 kg N₂O–N ha⁻¹ in this study, which fell well within the range reported in a meta-analysis for global croplands (0.3–16.8 kg N₂O–N ha⁻¹ y⁻¹) [43] and maize cropland in the Loess Plateau (0.65–3.52 kg N₂O–N ha⁻¹ y⁻¹) [44]. However, the mean annual cumulative N₂O emission in this study (1.38 kg N₂O–N ha⁻¹

y⁻¹) was significantly lower than that in the irrigated regions [41, 45, 46]. This may be because intensive water management, such as irrigation, was practiced in the irrigated regions, leading to an increase in the frequencies of alternate wetting and drying. Consequently, the C and N mineralization rates increased, which provided enough substrate for N₂O production [22, 37].

The background N₂O emission (0.80 kg N ha⁻¹ y⁻¹) fell within the range for various soil/climate regions and major cropping systems in P. R. China (0.1 to 3.67 kg N ha⁻¹ y⁻¹ with a mean of 1.5 kg N ha⁻¹ y⁻¹) [47], and accounted for 38.3–80.0% of the total N₂O emissions in the four N fertilization practices. A higher proportion of 81.8% was reported for a rain-fed wheat cropland in Western Australia [48], but its baseline N₂O emission (0.09 kg N ha⁻¹ y⁻¹) was approximately nine times lower than that in this study. The baseline N₂O emission was suggested to be resulting from nitrification and denitrification of soil indigenous N [47, 48]. The TN content (0.8 g kg⁻¹) and precipitation (550 mm) in this




Figure 5. Dynamics of soil nitrate content at 0–20 cm depths from fields with different nitrogen fertilization practices across fallow and maize growing season in 2013 and 2014. (FS, fallow season; MS, maize growing season).

study were higher than that in Western Australia (0.56 g kg^{-1} and 550 mm) [48]. In addition, in this study crop grew in hot and wet summer, but in cold and wet winter in Barton's study [48]. Thus, high TN content, temperature, and precipitation may contribute to the baseline N_2O emissions.

EF was $0.34 \pm 0.18\%$ in this study, which was within the range for fertilized uplands in P. R. China ($0.22\text{--}1.53\%$) [49], and the range for maize croplands in Canada ($0.03\text{--}1.45\%$) [50]. However, it was much lower than 1% suggested by IPCC [24] and 1.06% suggested by Davidson and Verchot [51]. Barton et al. [48] also reported a low EF (0.02%) in rain-fed croplands in semi-arid Southwestern Australia. Thus, the default value (1%) could cause an overestimation of direct N_2O emissions in the rain-fed cropland in the semi-arid regions, and multi-year measurements at multiple sites were needed to validate the emission factor for rain-fed croplands in semi-arid regions. EF was lower in rain-fed regions than in the irrigated regions, such as North China Plain ($0.61\text{--}0.77\%$), probably due to no irrigation and lower precipitation [52].

4.2 Factors controlling N_2O emissions

There was no significant relationship between soil temperature, WFPS, and N_2O emissions in this study, which was in agreement with other studies [8, 19]. However, N_2O emission was closely related to rainfall event ($>40 \text{ mm}$) [50, 53], and the peaks of N_2O emissions occurred after heavy rainfall events, such as on July 22, 2013 (120 mm), August 6 (44 mm) 2014, and September 30 (40.6 mm) 2014. After heavy rainfalls, the denitrification rate increased due to the formation of anaerobic environment [54]. In addition, the rates of C and N mineralization, as well as the cell lysis and intercellular solutes [55, 56] increased for several days following the wetting of dry soil [22, 37, 57]. As a result, soil mineral N increased, thus, providing enough substrate for N_2O production. For example, the $\text{NO}_3\text{--N}$ content after the heavy rainfall on July 20, 2013 was $20\text{--}176\%$ higher than that before rainfall. However, no N_2O peaks were observed after the




Figure 6. Dynamics of soil ammonium content at 0–20 cm depths from fields with different nitrogen fertilization practices across fallow and maize growing season in 2013 and 2014. (FS, fallow season; MS, maize growing season).

two rainfall events on September 18 (52 mm) and September 22 (40 mm) in 2013. This was probably due to the relatively low $\text{NO}_3\text{-N}$ (2.6–8.1 mg kg^{-1}) and $\text{NH}_4\text{-N}$ (3.8–8.1 mg kg^{-1}) contents, which limited the nitrification and denitrification rates [53, 58]. The cumulative N_2O emissions induced by rainfall accounted for 6.4 and 12.5% of the total annual N_2O emissions in 2013 and 2014, respectively.

Soil pH could also affect N cycling by directly affecting N mineralization and eventually N-containing gases [59]. Nitrogen mineralization linearly increased with soil pH, especially nitrification [60]. However, there was no significant relationship between pH and N mineralization in paddy waterlogged soils [61]. In calcareous soils with a high pH, ammonia volatilization was a loss pathway for fertilizer N and produced from soils under the condition of top dressing. Though the soil pH was 8.4 (calcareous soils), little ammonia volatilization appeared under the condition of buried fertilizer in this study [62].

There was a significantly positive correlation between N_2O emission and soil $\text{NO}_3\text{-N}$ across the four N fertilization practices, which was consistent with previous studies [44, 46, 48], indicating that the soil mineral N was a key variable controlling N_2O emissions [22, 42], especially within 10 days after N fertilization. It was noted that N_2O emissions within 10 days after N fertilization increased linearly with soil $\text{NO}_3\text{-N}$ content in both years 2013 and 2014 ($p < 0.05$). Liu et al. [42] also reported a linear relationship between N_2O emissions and soil mineral N content at the seasonal and annual scale. However, the slope was larger in 2014 (0.0093) than in 2013 (0.0058) (Fig. 7), probably due to the inter-annual variations in precipitation (523 mm in 2013 and 597 mm in 2014). The soil mean moisture within 10 days after N application was higher in 2014 (60%) than in 2013 (40%), which was conducive to the production of N_2O [8, 46, 63]. NI and SR also affected N_2O emissions by changing soil mineral N content. Figure 4 shows that the ammonium N content in all treatments reached a peak on April 27,




Figure 7. Relationship between mean soil NO₃-N at 0–20 cm depths and cumulative N₂O emissions within 10 days after N application.

2013, and it was higher in Opt + DCD treatment (45 mg kg⁻¹) than that in Opt treatment (42 mg kg⁻¹) because NI inhibited the first step of ammonium oxidation [31]. However, it was slightly lower in Opt + SR treatment (41 mg kg⁻¹), because SR fertilizers slowed down nutrient release rate through coating of the N fertilizers [11]. Thus, the highest nitrate N in Opt + DCD (98 mg kg⁻¹) and Opt + SR (151 mg kg⁻¹) treatments on May 23, 2013, was significantly lower compared with the Opt treatment (172 mg kg⁻¹) (Fig. 3). During the 2-year study period, the addition of NI significantly increased the ammonium N content and decreased the nitrate N content (Figs. 3 and 4). NI was effective in reducing N₂O emissions [8, 11, 64]. However, SR fertilizers had no such effect. In addition, Hu et al. found that the use of slow-release urea led to an increase in N₂O emissions (43%) during maize growing season and a decrease in N₂O emissions (33%) during wheat growing season [8]. This may be because high rainfall and irrigation during maize growing season accelerated SR hydrolysis [8], leading to N₂O emissions [9].

4.3 CH₄ uptakes and controlling factors

In this study, soils acted as a weak sink for atmospheric CH₄ by the oxidation of CH₄-oxidizing bacteria [7, 8, 45]. The annual CH₄ uptake ranged from 3.0 to 3.7 kg N CH₄-C ha⁻¹ with an average of 3.4 kg CH₄-C ha⁻¹, which was significantly higher than that in an eastern corn belt Alfisols in the USA [65] and a corn field in Indonesia [64]. N fertilizer was regarded as an inhibitor to CH₄ consumption in soils [21]. However, it had no significant effect on CH₄ uptakes in this study, which was also reported by Shi et al. [19]. There was no significant relationship between soil temperature, WFPS, and CH₄ uptakes, but CH₄ uptakes tended to increase from April to July and decreased slowly in winter, which was in line with a previous study [46].

4.4 Implications for food production and greenhouse gas emissions in the semi-arid Loess Plateau

Overuse of N fertilizers is common in P. R. China. The three optimized N fertilization practices saved 20% of N fertilizer compared with the conventional one (200 kg N ha⁻¹), without decreasing grain yield. Shi et al. also found no significant decrease

in grain yield with the reduction of N fertilizer from 200 to 185 or 186 kg N ha⁻¹ y⁻¹ in North China Plain [7]. After harvest, the nitrate accumulated in the 0–2 m soil profiles was reduced by 47.2, 48.5, and 45.5% in the three optimized N treatments as compared with the Con treatment, which minimized the risk of nitrate leaching.

The positive annual NGHG for all of the treatments (Tab. 1) indicated that N₂O emissions contributed significantly to the increase of NGHG in spite of CH₄ uptakes. Similar results were also reported for dry lands [7, 8, 66]. Compared to the Con treatment, NGHG was significantly reduced in the three optimized treatments due to their inhibitory effect on N₂O emissions. NGHGs ranged from 26 to 78 kg CO₂-eq Mg⁻¹ grain, which were comparable to that in maize systems in P. R. China (30–100 kg CO₂-eq Mg⁻¹ grain) [8]. The largest decrease of NGHGI (48.7% in 2013 and 58.7% in 2014) was observed in Opt + DCD treatment, followed by Opt + SR treatment (43.6% in 2013 and 39.7% in 2014). These results indicated that Opt + DCD and Opt + SR treatments could significantly decrease N₂O emissions from soils while maintaining a high maize yield in the rain-fed regions of Loess Plateau. The spring maize is mainly planted in northern China with a plant area of seven million hectares [19], thus, the application of the most effective N fertilization practice, Opt + DCD, could save 0.35 million Mg N y⁻¹ and 6.5 million kg N₂O-N y⁻¹.

The economic costs and benefits of these new N fertilization practices also need to be taken into account. The slow release urea used in this study was expensive relative to urea because of the complex manufacturing technology, and the NI used in this study was chemically pure, and thus, also expensive. Consequently, although NI has the potential to significantly decrease GHG emissions, farmers may be less willing to use it because of extra economic costs. A promising alternative solution is to add NI directly in the production of urea. In addition, subsidies or intervention policies from local government are also necessary for the application of these new N fertilization practices.

5 Concluding remarks

This 2-year field study investigated the effects of optimized N fertilization practices on GHG emissions in a rain-fed spring maize cropland. Soil acted as a small sink for atmospheric CH₄, and N fertilization practices had no significant effect on CH₄ uptakes. Soil acted as a source of N₂O, and N fertilization practices significantly influenced N₂O emissions. Soil temperature and moisture had no significant effects on N₂O emissions, but were closely related to fertilization and rainfall events (>40 mm). The background N₂O emissions and EFs in the rain-fed regions of Loess Plateau were much lower than the default values suggested by IPCC.

The three optimized N fertilization practices saved 20% of N fertilizer compared with the conventional treatment, but without a significant decrease in grain yield; and decreased the annual N₂O emissions, NGHG, and NGHGI. NI decreased cumulative N₂O emissions by about 72% within 10 days after N fertilization. Compared to the conventional treatment, NI significantly reduced annual N₂O emissions, NGHG, and NGHGI by 48, 56, and 54% on average, respectively. Moreover, NI decreased the accumulated nitrate in the 0–2 m soil profiles after harvest and minimized the risk of nitrate leaching. Therefore, Opt + DCD is the most effective strategy to mitigate N₂O emissions from the maize systems in the rain-fed regions of Loess Plateau. Reducing the cost of urea with NI was crucial for its widespread applications in agriculture.

Acknowledgments

This work is supported by the Special Fund for the Agricultural Profession (201103039), and the “Strategic Priority Research Program-Climate Change: Carbon Budget and Related Issues” of the Chinese Academy of Sciences (XDA05050504).

The authors have declared no conflict of interest.

References

- [1] D. S. Reay, E. A. Davidson, K. A. Smith, P. Smith, J. M. Melillo, F. Dentener, P. J. Crutzen, Global Agriculture and Nitrous Oxide Emissions, *Nat. Clim. Change* **2012**, *2* (6), 410–416.
- [2] P. Smith, D. Martino, Z. C. Cai, D. Gwary, H. Janzen, P. Kumar, B. McCarl, et al., Agriculture, in *Climate Change 2007: Mitigation* (Eds.: B. Metz, O. R. Davidson, P. R. Bosch, R. Dave, L. A. Meyer), Contribution of Working Group III to the fourth assessment report of the intergovernmental panel on climate change, Cambridge University Press, Cambridge **2007**, pp. 497–540.
- [3] L. Bouwman, J. S. Daniel, E. A. Davidson, C. de Klein, E. Holland, X. Ju, D. Kanter, et al., *Drawing Down N₂O to Protect Climate and the Ozone Layer*, A UNEP Synthesis Report, UNEP, Nairobi, Kenya **2013**.
- [4] A. R. Mosier, A. D. Halvorson, C. A. Reule, X. J. Liu, Net Global Warming Potential and Greenhouse Gas Intensity in Irrigated Cropping Systems in Northeastern Colorado, *J. Environ. Qual.* **2006**, *35* (4), 1584–1598.
- [5] G. P. Robertson, P. M. Vitousek, Nitrogen in Agriculture: Balancing the Cost of an Essential Resource, *Annu. Rev. Environ. Resour.* **2009**, *34*, 97–125.
- [6] U. M. Sainju, J. L. Barsotti, J. Wang, Net Global Warming Potential and Greenhouse Gas Intensity Affected by Cropping Sequence and Nitrogen Fertilization, *Soil Sci. Soc. Am. J.* **2014**, *78* (1), 248–261.
- [7] Y. F. Shi, W. L. Wu, F. Q. Meng, L. Zheng, D. P. Wang, H. Ye, G. W. Ding, Nitrous Oxide and Methane Fluxes During the Maize Season Under Optimized Management in Intensive Farming Systems of the North China Plain, *Pedosphere* **2014**, *24* (4), 487–497.
- [8] X. K. Hu, F. Su, X. T. Ju, B. Gao, O. Oenema, P. Christie, B. X. Huang, et al., Greenhouse Gas Emissions From a Wheat-Maize Double Cropping System With Different Nitrogen Fertilization Regimes, *Environ. Pollut.* **2013**, *176*, 198–207.
- [9] H. Akiyama, X. Yan, K. Yagi, Evaluation of Effectiveness of Enhanced-Efficiency Fertilizers as Mitigation Options for N₂O and NO Emissions From Agricultural Soils: Meta-Analysis, *Global Change Biol.* **2010**, *16* (6), 1837–1846.
- [10] J. Y. Jiang, Z. H. Hu, W. J. Sun, Y. H. Huang, Nitrous Oxide Emissions From Chinese Cropland Fertilized With a Range of Slow-Release Nitrogen Compounds, *Agric. Ecosyst. Environ.* **2010**, *135* (3), 216–225.
- [11] E. Guertal, Slow-Release Nitrogen Fertilizers in Vegetable Production: A Review, *HortTechnology* **2009**, *19* (1), 16–19.
- [12] J. G. Gu, X. H. Zheng, Y. H. Wang, W. X. Ding, B. Zhu, X. Chen, Y. Y. Wang, et al., Regulatory Effects of Soil Properties on Background N₂O Emissions From Agricultural Soils in China, *Plant Soil* **2007**, *295* (1–2), 53–65.
- [13] H. Di, K. Cameron, The Use of a Nitrification Inhibitor, Dicyandiamide (DCD), to Decrease Nitrate Leaching and Nitrous Oxide Emissions in a Simulated Grazed and Irrigated Grassland, *Soil Use Manage.* **2002**, *18* (4), 395–403.
- [14] C. de Klein, Potential mitigation options for reducing methane and nitrous oxide emissions from dairy farms, in *Dairy Farm Soil Management* (Eds.: L. D. Currie, P. Loganathan), Occasional report No. 15, Fertilizer and Lime Research Centre, Massey University, Palmerston North, New Zealand **2002**, pp. 233–246.
- [15] K. E. Dobbie, K. A. Smith, Nitrous Oxide Emission Factors for Agricultural Soils in Great Britain: The Impact of Soil Water-Filled Pore Space and Other Controlling Variables, *Global Change Biol.* **2003**, *9* (2), 204–218.
- [16] C. De Klein, L. Smith, R. Monaghan, Restricted Autumn Grazing to Reduce Nitrous Oxide Emissions From Dairy Pastures in Southland, New Zealand, *Agric. Ecosyst. Environ.* **2006**, *112* (2), 192–199.
- [17] A. R. Mosier, J. A. Delgado, M. Keller, Methane and Nitrous Oxide Fluxes in an Acid Oxisol in Western Puerto Rico: Effects of Tillage, Liming and Fertilization, *Soil Biol. Biochem.* **1998**, *30* (14), 2087–2098.
- [18] K. Bronson, A. Mosier, Suppression of Methane Oxidation in Aerobic Soil by Nitrogen Fertilizers, Nitrification Inhibitors, and Urease Inhibitors, *Biol. Fertil. Soils* **1994**, *17* (4), 263–268.
- [19] Y. F. Shi, W. L. Wu, F. Q. Meng, Z. H. Zhang, L. Zheng, D. P. Wang, Integrated Management Practices Significantly Affect N₂O Emissions and Wheat-Maize Production at Field Scale in the North China Plain, *Nutr. Cycl. Agroecosyst.* **2013**, *95* (2), 203–218.
- [20] J. Jing, F. Zhang, Z. Rengel, J. Shen, Localized Fertilization With P Plus N Elicits an Ammonium-Dependent Enhancement of Maize Root Growth and Nutrient Uptake, *Field Crop Res.* **2012**, *133*, 176–185.
- [21] N. C. Turner, S. Asseng, Productivity, Sustainability, and Rainfall-Use Efficiency in Australian Rainfed Mediterranean Agricultural Systems, *Crop Pasture Sci.* **2005**, *56* (11), 1123–1136.
- [22] C. Snyder, T. Bruulsema, T. Jensen, P. Fixen, Review of Greenhouse Gas Emissions From Crop Production Systems and Fertilizer Management Effects, *Agric. Ecosyst. Environ.* **2009**, *133* (3), 247–266.
- [23] J. Hoben, R. Gehl, N. Millar, P. Grace, G. Robertson, Nonlinear Nitrous Oxide (N₂O) Response to Nitrogen Fertilizer in On-Farm Corn Crops of the US Midwest, *Global Change Biol.* **2011**, *17* (2), 1140–1152.
- [24] IPCC, *Guidelines for National Greenhouse Gas Inventories*, IGES, Hayama, Japan **2006**.
- [25] A. Bouwman, Direct Emission of Nitrous Oxide From Agricultural Soils, *Nutr. Cycl. Agroecosyst.* **1996**, *46* (1), 53–70.
- [26] E. G. Gregorich, P. Rochette, A. J. Vanden Bygaart, D. A. Angers, Greenhouse Gas Contributions of Agricultural Soils and Potential Mitigation Practices in Eastern Canada, *Soil Till. Res.* **2005**, *83* (1), 53–72.
- [27] S. L. Guo, J. S. Wu, K. Coleman, H. H. Zhu, Y. Li, W. Z. Liu, Soil Organic Carbon Dynamics in a Dryland Cereal Cropping System of the Loess Plateau Under Long-Term Nitrogen Fertilizer Applications, *Plant Soil* **2011**, *353* (1–2), 321–332.
- [28] M. B. Huang, T. H. Dang, J. Gallichand, M. Goulet, Effect of Increased Fertilizer Applications to Wheat Crop on Soil-Water Depletion in the Loess Plateau, China, *Agric. Water Manage.* **2003**, *58* (3), 267–278.
- [29] T. L. Fan, B. Stewart, W. A. Payne, W. Yong, J. J. Luo, Y. F. Gao, Long-Term Fertilizer and Water Availability Effects on Cereal Yield and Soil Chemical Properties in Northwest China, *Soil Sci. Soc. Am. J.* **2005**, *69* (3), 842–855.
- [30] X. P. Deng, L. Shan, H. P. Zhang, N. C. Turner, Improving Agricultural Water Use Efficiency in Arid and Semiarid Areas of China, *Agric. Water Manage.* **2006**, *80* (1), 23–40.
- [31] J. L. Liu, L. D. Bu, L. Zhu, S. S. Luo, X. P. Chen, S. Q. Li, R. L. Hill, et al., Nitrogen Fertilization Effects on Nitrogen Balance and Use Efficiency for Film-Mulched Maize in a Semiarid Region, *Acta Agric. Scand. Sect. B* **2013**, *63* (7), 612–622.
- [32] S. L. Guo, H. H. Zhu, T. H. Dang, J. S. Wu, W. Z. Liu, M. D. Hao, Y. Li, et al., Winter Wheat Grain Yield Associated With Precipitation Distribution Under Long-Term Nitrogen Fertilization in the Semiarid Loess Plateau in China, *Geoderma* **2012**, *189*, 442–450.
- [33] S. X. Li, Z. H. Wang, S. Malhi, S. Q. Li, Y. J. Gao, X. H. Tian, Nutrient and Water Management Effects on Crop Production, and Nutrient and Water Use Efficiency in Dryland Areas of China, *Adv. Agron.* **2009**, *102*, 223–265.
- [34] M. K. Firestone, E. A. Davidson, M. Andreae, D. Schimel, Microbiological basis of NO and N₂O production and consumption in soil, in *Exchange of Trace Gases Between Terrestrial Ecosystems and the Atmosphere* (Eds.: M. O. Andreae, D. S. Schimel), John Wiley & Sons, Chichester, UK **1989**, pp. 7–21.
- [35] X. R. Wei, M. D. Hao, X. H. Xue, P. Shi, R. Horton, A. Wang, Y. F. Zang, Nitrous Oxide Emission From Highland Winter Wheat Field After Long-Term Fertilization, *Biogeosciences* **2010**, *7* (10), 3301–3310.
- [36] R. X. Shao, L. Deng, Q. H. Yang, Z. P. Shangguan, Nitrogen Fertilization Increase Soil Carbon Dioxide Efflux of Winter Wheat Field: A Case Study in Northwest China, *Soil Tillage Res.* **2014**, *143*, 164–171.

- [37] Y. Zhang, H. Zhang, Y. B. Qi, Y. J. Hao, Y. F. Bai, Soil Nitrate Nitrogen Distribution and Transportation Under Several Vegetations on the Reconstructed Sloping Land in the Loess Plateau, *J. Agro-Environ. Sci.* **2012**, *31* (1), 155–160.
- [38] S. Hao, Y. S. Yang, Z. J. Yang, G. S. Chen, J. S. Xie, J. F. Guo, S. Zou, The Dynamic Response of Soil Respiration to Land-Use Changes in Subtropical China, *Global Change Biol.* **2010**, *16* (3), 1107–1121.
- [39] M. B. Huang, M. D. Shao, L. Zhang, Y. S. Li, Water Use Efficiency and Sustainability of Different Long-Term Crop Rotation Systems in the Loess Plateau of China, *Soil Tillage Res.* **2003**, *72* (1), 95–104.
- [40] X. H. Zheng, B. L. Mei, Y. H. Wang, B. H. Xie, Y. S. Wang, H. B. Dong, H. Xu, et al., Quantification of N₂O Fluxes From Soil-Plant Systems May Be Biased by the Applied Gas Chromatograph Methodology, *Plant Soil* **2008**, *311* (1–2), 211–234.
- [41] C. Liu, K. Wang, X. Zheng, Responses of N₂O and CH₄ Fluxes to Fertilizer Nitrogen Addition Rates in an Irrigated Wheat-Maize Cropping System in Northern China, *Biogeosciences* **2012**, *9* (2), 839–850.
- [42] J. L. Liu, L. B. Zhu, S. S. Luo, L. T. Bu, X. P. Chen, S. C. Yue, S. Q. Li, Response of Nitrous Oxide Emission to Soil Mulching and Nitrogen Fertilization in Semi-Arid Farmland, *Agric. Ecosyst. Environ.* **2014**, *188*, 20–28.
- [43] P. Forster, V. Ramaswamy, P. Artaxo, T. Bernsten, R. Betts, D. W. Fahey, J. Hayward, et al., Changes in atmospheric constituents and in radiative forcing, Chapter 2, in *Climate Change 2007, The Physical Science Basis, Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* (Eds.: S. Solomon, D. Qin, M. Manning, Z. Chen, M. Marquis, K. B. Averyt, M. Tignor, et al.), Cambridge University Press, Cambridge, UK and New York, NY, USA **2007**.
- [44] E. Stehfest, L. Bouwman, N₂O and NO Emission From Agricultural Fields and Soils Under Natural Vegetation: Summarizing Available Measurement Data and Modeling of Global Annual Emissions, *Nutr. Cycl. Agroecosyst.* **2006**, *74* (3), 207–228.
- [45] B. Gao, X. T. Ju, F. Su, Q. F. Meng, O. Oenema, P. Christie, X. P. Chen, et al., Nitrous Oxide and Methane Emissions From Optimized and Alternative Cereal Cropping Systems on the North China Plain: A Two-Year Field Study, *Sci. Total Environ.* **2014**, *472*, 112–124.
- [46] S. Riya, J. Min, S. Zhou, W. M. Shi, M. Hosomi, Short-Term Responses of Nitrous Oxide Emissions and Concentration Profiles to Fertilization and Irrigation in Greenhouse Vegetable Cultivation, *Pedosphere* **2012**, *22* (6), 764–775.
- [47] S. H. Yang, S. Z. Peng, J. Z. Xu, Y. F. Luo, D. X. Li, Methane and Nitrous Oxide Emissions From Paddy Field as Affected by Water-Saving Irrigation, *Phys. Chem. Earth* **2012**, *53–54*, 30–37.
- [48] L. Barton, R. Kiese, D. Gatter, K. Butterbach-Bahl, R. Buck, C. Hinz, D. V. Murphy, Nitrous Oxide Emissions From a Cropped Soil in a Semi-Arid Climate, *Global Change Biol.* **2008**, *14* (1), 177–192.
- [49] R. Lemke, R. Izaurrealde, M. Nyborg, Seasonal Distribution of Nitrous Oxide Emissions From Soils in the Parkland Region, *Soil Sci. Soc. Am. J.* **1998**, *62* (5), 1320–1326.
- [50] B. L. Ma, T. Y. Wu, N. Tremblay, W. Deen, M. J. Morrison, N. B. McLaughlin, E. G. Gregorich, et al., Nitrous Oxide Fluxes From Corn Fields: On-Farm Assessment of the Amount and Timing of Nitrogen Fertilizer, *Global Change Biol.* **2010**, *16* (1), 156–170.
- [51] E. A. Davidson, L. V. Verchot, Testing the Hole In-the-Pipe Model of Nitric and Nitrous Oxide Emissions From Soils Using the TRAGNET Database, *Global Biogeochem. Cycles* **2000**, *14* (4), 1035–1043.
- [52] W. X. Ding, Y. Cai, Z. C. Cai, K. Yagi, X. H. Zheng, Nitrous Oxide Emissions From an Intensively Cultivated Maize-Wheat Rotation Soil in the North China Plain, *Sci. Total Environ.* **2007**, *373* (2–3), 501–511.
- [53] C. Y. Liu, K. Wang, S. X. Meng, X. H. Zheng, Z. X. Zhou, S. H. Han, D. L. Chen, et al., Effects of Irrigation, Fertilization and Crop Straw Management on Nitrous Oxide and Nitric Oxide Emissions From a Wheat-Maize Rotation Field in Northern China, *Agric. Ecosyst. Environ.* **2011**, *140* (1–2), 226–233.
- [54] X. T. Ju, X. Lu, Z. L. Gao, X. P. Chen, F. Su, M. Kogge, V. Romheld, et al., Processes and Factors Controlling N₂O Production in an Intensively Managed Low Carbon Calcareous Soil Under Sub-Humid Monsoon Conditions, *Environ. Pollut.* **2011**, *159* (4), 1007–1016.
- [55] M. Xing, W. G. Liu, Z. F. Wang, J. Hu, Relationship of Nitrate Isotopic Character to Population Density in the Loess Plateau of Northwest China, *Appl. Geochem.* **2013**, *35*, 110–119.
- [56] A. Bouwman, L. Boumans, N. Batjes, Emissions of N₂O and NO From Fertilized Fields: Summary of Available Measurement Data, *Global Biogeochem. Cycles* **2002**, *16* (4), 6–16–13.
- [57] J. Delgado, A. Mosier, Mitigation Alternatives to Decrease Nitrous Oxides Emissions and Urea-Nitrogen Loss and Their Effect on Methane Flux, *J. Environ. Qual.* **1996**, *25* (5), 1105–1111.
- [58] L. Horváth, B. Grosz, A. Machon, Z. Tuba, Z. Nagy, S. Czóbel, J. Balogh, et al., Estimation of Nitrous Oxide Emission From Hungarian Semi-Arid Sandy and Loess Grasslands; Effect of Soil Parameters, Grazing, Irrigation and Use of Fertilizer, *Agric. Ecosyst. Environ.* **2010**, *139* (1), 255–263.
- [59] D. Curtin, C. A. Campbell, A. Jalil, Effects of Acidity on Mineralization: pH-Dependence of Organic Matter Mineralization in Weakly Acidic Soils, *Soil Biol. Biochem.* **1998**, *30* (1), 57–64.
- [60] D. J. Mladenoff, Dynamics of Nitrogen Mineralization and Nitrification in Hemlock and Hardwood Treefall Gaps, *Ecology* **1987**, *68* (5), 1171–1180.
- [61] K. L. Sahrawat, Mineralization of Soil Organic Nitrogen Under Waterlogged Conditions in Relation to Other Properties of Tropical Rice Soils, *Soil Res.* **1983**, *21* (2), 133–138.
- [62] X. T. Ju, G. X. Xing, X. P. Chen, S. L. Zhang, L. J. Zhang, X. J. Liu, F. S. Zhang, Reducing Environmental Risk by Improving N Management in Intensive Chinese Agricultural Systems, *Proc. Natl. Acad. Sci. USA* **2009**, *106* (9), 3041–3046.
- [63] Y. Q. Wang, X. C. Zhang, C. Q. Huang, Spatial Variability of Soil Total Nitrogen and Soil Total Phosphorus Under Different Land Uses in a Small Watershed on the Loess Plateau, China, *Geoderma* **2009**, *150* (1), 141–149.
- [64] K. Smith, T. Ball, F. Conen, K. Dobbie, J. Massheder, A. Rey, Exchange of Greenhouse Gases Between Soil and Atmosphere: Interactions of Soil Physical Factors and Biological Processes, *Eur. J. Soil Sci.* **2003**, *54* (4), 779–791.
- [65] D. R. Smith, G. Hernandez-Ramirez, S. D. Armstrong, D. L. Bucholtz, D. E. Stott, Fertilizer and Tillage Management Impacts on Non-Carbon-Dioxide Greenhouse Gas Emissions, *Soil Sci. Soc. Am. J.* **2011**, *75* (3), 1070–1082.
- [66] R. S. Jassal, T. A. Black, R. Roy, G. Ethier, Effect of Nitrogen Fertilization on Soil CH₄ and N₂O Fluxes, and Soil and Bole Respiration, *Geoderma* **2011**, *162* (1–2), 182–186.